PHARMACOPŒIA

OF THE

LONDON HOSPITAL.

M
25329

QV740
.BE5
1908.
L84P

2

1908.

M25329

22500288319

75519

PHARMACOPŒIA

OF THE

LONDON HOSPITAL,

COMPILED

Under the direction of a Committee
appointed by

THE LONDON HOSPITAL MEDICAL COUNCIL.

London:
SCHOOL-PRESS, Rupert Street, London, E.

Price 1/6.

1908

24 880 135

12867

M25329

WELLCOME INSTITUTE LIBRARY

Coll.	WelMOmec
Coll. No.	QV740 .BE5 1908 L84p

TABLE OF CONTENTS.

	PAGE.
ANTIDOTE TABLE	5
BALNEA...	31
COLLODIUM	32
COLLUNARIA	33
CONFECTIONES...	34
DENTIFRICIA	34
EMPLASTRUM	35
EMULSIONES	35
ENEMATA	35
EXTRACTA	37
FOTUS	37
GARGARISMATA	38
GLYCERINA	40
GUTTÆ	40
HAUSTUS	43
INJECTIONES	44
LINCTUS	47
LINIMENTA	47
LIQUORES	48
LOTIONES	49
MISTURÆ	53
NEBULÆ	63

PAGE.

PASTÆ 65
PIGMENTA 65
PILULÆ 67
POTUS 68
PULVERES 68
SUPPOSITORIA 70
SYRUPI 71
TABELLÆ 71
TROCHISCUS 77
UNGUENTA 77
VAPORES 82
BEEF JUICE 84
,, TEA 84
EGG WHITE SOLUTION 85
NORMAL SALINE 85
PEPTONISED MILK 85
DIET WITH MINIMUM QUANTITY OF CARBO-HYDRATES 86
INFANT FEEDING 87
DIRECTIONS FOR THE CARE OF RICKETY CHILDREN 89
THERMOMETRIC TABLE 90
WEIGHTS AND MEASURES 91
POSOLOGICAL TABLE 95

ANTIDOTE TABLE.

When an Emetic is ordered, one of the following, unless a special Emetic is mentioned, should be given:—

(*a*) *Mustard.*—A tablespoonful in two-thirds of a tumbler of luke-warm water.

(*b*) *Zinc Sulphate.*—30 grains in half a tumbler of luke-warm water.

(*c*) *Powdered Ipecacuanha Root.*—10 grains in half a tumbler of luke-warm water, repeat if necessary.

(*d*) *Warm Water.*—Given copiously alone or after one of the above.

(*e*) Mechanical stimulation of fauces and back of pharynx with finger or feather.

(*f*) *Apomorphine Hydrochloride.*—$\frac{1}{20}$ to $\frac{1}{4}$ grain, hypodermically.

When Demulcents are ordered, one or more of the following may be given:—

(*a*) *Milk.*

(*b*) *Linseed Tea.*

(*c*) *Barley Water.*

When Stimulants are ordered, one or more of the following may be given:—

(*a*) *Brandy or Whisky.*—1 to 2 ounces with 4 ounces of warm water by mouth, or ½ to 1 ounce with 2 ounces of warm water by rectum; or 1 to 2 drachms hypodermically.

(*b*) *Mixture of Ammonia and Ether.*—1 ounce.

(*c*) *Sal Volatile.*—2 drachms with 2 ounces of water.

(*d*) *Ether.*—30 to 60 minims hypodermically.

Aconite.

Immediate Treatment.—Stomach Tube. Solution of Atropine Sulphate, 4 minims, by mouth; or Atropine Sulphate, $\frac{1}{30}$ grain, hypodermically; or Solution of Adrenalin, 10 minims, hypodermically.

Later.—Brandy. Warmth. Friction. Artificial respiration.

Alcohol.

Immediate Treatment.—Stomach Tube. Cold Affusions to head.

Later.—Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Solution of Strychnine Hydrochloride, 5 minims, ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically. Repeat if necessary. Warmth.

Ammonia.

Stomach Tube, with care. Vinegar or Diluted Acetic Acid, or Lemon Juice, 1 ounce in 3 ounces of Water. Olive Oil. Tracheotomy if necessary.

Antimony.

Stomach Tube or Emetic, if required. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water, to be repeated if vomited. Strong Tea or Coffee, ½ pint. Milk. Stimulants. Hot Bottles and Blankets. Friction.

Later.—Morphine Sulphate, ¼ grain, hypodermically. Repeat in 1 hour if necessary.

Aqua Fortis.

Immediate Treatment.—Soap and Water. Calcined Magnesia, Chalk or Whiting, ½ ounce in ½ pint of Water or Milk; or Sodium Bicarbonate, 2 drachms in 5 ounces of Water.

Later.—Demulcents. Olive Oil. White of Egg and Water. Morphine Sulphate, ¼ grain, hypodermically.

Arsenic or Arsenious Acid.

Immediate Treatment.—Stomach Tube or Emetic. Ferric Hydrate (prepared by mixing Solution of Ferric Chloride, ½ ounce in half a tumbler of Water, with an equal quantity of Saturated Solution of Washing Soda). Collect precipitate on a handkerchief and mix with half a tumbler of Hot Water. Repeat frequently. (Solution of Ammonia may be used instead of Washing Soda).

Atropine.

Stomach Tube or Emetic. Brandy. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Pilocarpine Nitrate, ½ grain, hypodermically, to be repeated in 15 minutes if pulse improve. Flicking. Pinching. Warm and Cold Douche. Faradism. Hot Bottles and Blankets. Artificial respiration. Friction.

Barium, Salts of.

Stomach Tube or Emetic. Sodium Sulphate, 1 ounce in 5 ounces of Water; or Magnesium Sulphate, ½ ounce in 5 ounces of Water; or Alum, 1 drachm in 5 ounces of Water. Stimulants. Hot Bottles and Blankets.

Later.—Morphine Sulphate, ¼ grain, hypodermically.

Battley's Solution (*See Opium*).

Belladonna.

Stomach Tube or Emetic. Brandy. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Pilocarpine Nitrate, ½ grain, hypodermically, to be repeated in 15 minutes if pulse improve. Flicking. Pinching. Warm and Cold Douche. Faradism. Hot Bottles and Blankets Artificial respiration. Friction.

Benzol.

Immediate Treatment.—Stomach Tube. Emetic.

Later.—Brandy. Inhalation of Ammonia. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically.

Bichromate of Potassium.

Stomach Tube or Emetic. Chalk Mixture, 8 ounces; or Chalk, ½ ounce in ½ pint of Milk. Hot Bottles and Blankets.

Burnett's Disinfecting Fluid (Zinc Chloride).

White of Egg and Water. Washing Soda in dilute solution. Demulcents. Strong Tea, ½ pint. Hot Fomentations. Morphine Sulphate, ¼ grain, hypodermically.

Calabar Bean (Eserine).

Emetics. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth every 15 minutes until pupils dilate. Chloral Hydrate, 10 grains in 2 ounces of Water, by mouth or rectum every 15 minutes for 4 doses. Stimulants. Artificial respiration. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{120}$ grain of Strychnine Hydrochloride), hypodermically.

Camphor.

Immediate Treatment. — Stomach Tube and Emetic.

Later.—Inhalation of Ammonia. Subcutaneous injection of Brandy. Hot Fomentations.

Cantharides.

Immediate Treatment.—Stomach Tube or Emetic.

Later.—Demulcents. Morphine Sulphate, ¼ grain, hypodermically.

Carbolic Acid (Phenol).

Stomach Tube; passed under Anæsthetic if necessary. Sodium Sulphate, 1 ounce in 3 ounces of Water; or Saccharated Lime Water, 1 ounce in 3 ounces of Water.

Later.—Demulcents. Olive Oil. Brandy, subcutaneously or by enema. Warmth.

Carbonic Acid Gas.

Fresh air. Artificial respiration. Inhalation of Ammonia. Stimulants. Hot Bottles and Blankets. Inhalation of Oxygen. Faradism. Cold Douche.

Carbon Monoxide.

Fresh air. Artificial respiration. Bleeding and Intravenous injection of Warm Saline Solution. Inhalation of Oxygen. Stimulants. Hot Bottles and Blankets. Friction. Faradism.

Carbon Tetrachloride (*See Ether*).

Caustic Potash and Soda.

Stomach Tube with care. Vinegar or Diluted Acetic Acid, or Lemon Juice, 1 ounce in 5 ounces of Water. Olive Oil.

Cesspool Gas.

Fresh air. Artificial respiration. Stimulants. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Hot Bottles and Blankets. Faradism.

Chloral Hydrate.

Immediate Treatment.—Prevent sleep. Stomach Tube and Emetic. Hot Blankets. Hot Bottle to feet.

Later.—Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Artificial respiration. Faradism. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically. Capsule of Amyl Nitrite for inhalation. Inhalation of Oxygen.

Chlorate of Potassium.

Emetic. Wash out Stomach.

Chlorodyne.

Stomach Tube or Emetic. Wash out Stomach at half hour intervals with Solution of Potassium Permanganate, diluted with five times the quantity of Warm Water, leaving about 5 ounces of the Diluted Solution in the Stomach. Mixture of Ammonia and Ether, 1 ounce; or Ether, 60 minims, hypodermically. Pinching. Flapping. Solution of Ammonia or Smelling Salts to nostrils. Faradism. Make patient walk about. Solution of Atropine Sulphate, 3 minims ($\frac{3}{100}$ grain of Atropine Sulphate), hypodermically, to be repeated in half an hour; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth or rectum. Capsule of Amyl Nitrite for inhalation. Artificial respiration.

Chloroform (*Inhaled*).

Insert finger into mouth and pharynx to make sure that there is no obstruction, *e.g.*, artificial teeth. Extend the head and push the jaw forward. Pull out tongue with forceps. Loosen clothes. Fresh air. Artificial respiration. Raise foot of table that the head may be lower than the rest of the body. Flapping with wet towels. Stimulants. Intravenous injection of Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), to be washed through with Normal Saline.

Chloroform (*Swallowed*).

Stomach Tube or Emetic. Olive Oil. Stimulants, hypodermically or by rectum. Flapping with wet towels. Faradism. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Capsule of Amyl Nitrite for inhalation.

Coal Gas.

Fresh air. Artificial respiration. Stimulants. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Inhalation of Oxygen. Inhalation of Ammonia. Transfusion.

Cocaine.

Stomach Tube, if swallowed. Stimulants. Capsule of Amyl Nitrite for inhalation.

Colchicum.

Emetic. Olive Oil or any Demulcent. Stimulants.

Copper, Salts of.

Immediate Treatment. — White of Egg and Water. Milk. Stomach Tube if vomiting has not occurred.

Later.—Demulcents. Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically, and Hot Fomentations if much pain.

Corrosive Sublimate.

Stomach Tube, with great care. Flour and Water; or Arrowroot and Water; or White of Egg and Water, *ad lib.* Leave in Stomach White of Egg and Water (1 egg = 4 grains of Corrosive Sublimate). Stimulants. Hot Bottles and Blankets.

Later.—Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically.

Creosote.

Stomach Tube. Sodium Sulphate, 1 ounce in 3 ounces of Water; or Saccharated Lime Water, 1 ounce in 3 ounces of Water.

Later.—Demulcents. Olive Oil. Brandy, subcutaneously or by enema. Warmth.

Croton Oil.

Stomach Tube or Emetic. Stimulants. Morphine Sulphate, $\frac{1}{2}$ grain, hypodermically; or Tincture of Opium, 30 minims in 1 ounce of Water, by mouth or rectum; or Morphine Suppository. Hot Bottles and Blankets. Friction. Poultices or Hot Fomentations to abdomen.

Cyanide of Potassium.

Stomach Tube. Wash out Stomach with 2% Solution of Copper Sulphate. Stimulants, hypodermically. Faradism. Inhalation of Ammonia. Artificial respiration. Fresh air. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth or rectum. Alternate Warm and Cold Douches. Friction.

Deadly Nightshade.

Stomach Tube or Emetic. Brandy. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Pilocarpine Nitrate, $\frac{1}{2}$ grain, hypodermically, to be repeated in 15 minutes if pulse improve. Flicking. Pinching. Warm and Cold Douche. Faradism. Hot Bottles and Blankets. Artificial respiration. Friction.

Digitalis (Foxglove).

Emetics. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water. Stimulants. Recumbent position. Solution of Atropine Sulphate, 1 minim ($\frac{1}{100}$ grain of Atropine Sulphate), hypodermically; and Solution of Trinitrin, 2 minims, by mouth.

Eserine.

Emetics. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate, hypodermically; or Tincture of Belladonna, 15 minims, by mouth every 15 minutes until pupils dilate. Chloral Hydrate, 10 grains in 2 ounces of Water, by mouth or rectum every 15 minutes for 4 doses. Stimulants. Artificial respiration. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically.

Ether (*Inhaled*).

Insert finger into mouth and pharynx to make sure that there is no obstruction, *e.g.*, artificial teeth. Extend the head and push the jaw forward. Pull tongue out with forceps. Loosen clothes. Fresh air. Artificial respiration. Raise foot of table that the head may be lower than the rest of the body. Flapping with wet towels. Intravenous injection of Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), to be washed through with Normal Saline.

Ether (*Swallowed*).

Stomach Tube or Emetic. Olive Oil. Stimulants, hypodermically or by rectum. Flapping with wet towels. Faradism. Capsule of Amyl Nitrite for inhalation. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically.

Foxglove (Digitalis).

Emetics. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water. Stimulants. Recumbent position. Solution of Atropine Sulphate, 1 minim ($\frac{1}{100}$ grain of Atropine Sulphate), hypodermically; and Solution of Trinitrin, 2 minims, by mouth.

Fungi.

Emetics. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically. Stimulants.

Henbane.

Stomach Tube or Emetic. Brandy. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Pilocarpine Nitrate, $\frac{1}{2}$ grain, hypodermically, to be repeated in 15 minutes if pulse improve. Flicking. Pinching. Warm and Cold Douche. Faradism. Hot Bottles and Blankets. Artificial respiration. Friction.

Hydrochloric Acid (Muriatic Acid, Spirits of Salts).

Introduce Water and use Stomach Tube with care. Sodium Bicarbonate, 2 drachms in 5 ounces of Water; or Soap and Water; or Chalk or Calcined Magnesia, $\frac{1}{2}$ ounce in $\frac{1}{2}$ pint of Water or Milk; or Solution of Ammonia or Sal Volatile, freely diluted; or White of Egg and Water; or Milk.

Later. — Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically, for pain, and Ice for thirst.

Hydrocyanic Acid.

Immediate Treatment.—Dash Cold Water over Head and Chest. Wash out Stomach with 2% Solution of Copper Sulphate. Emetic. Inhalation of Ammonia. Artificial respiration. Solution of Atropine Sulphate 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically.

Iodine.

Immediate Treatment. — Stomach Tube and Emetic. Wash out Stomach with Diluted Solution of Potash.

Later.—Starch and Water. Demulcents. Capsule of Amyl Nitrite for inhalation. Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically, and Hot Fomentations for pain.

Lead and its Compounds.

Stomach Tube or Emetic. Magnesium Sulphate, $\frac{1}{2}$ ounce in 5 ounces of Water; or Sodium Sulphate, 1 ounce in 5 ounces of Water; or Diluted Sulphuric Acid, 30 minims in $\frac{1}{2}$ pint of Water. Milk; or White of Egg and Water. Poultices to Abdomen. Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically.

Liniment of Aconite (*See Aconite*).

Liniment of Aconite, Belladonna and Chloroform—*A.B.C.* (*See Aconite, Belladonna and Chloroform*).

Liniment of Ammonia (*See Ammonia*).

Liniment of Belladonna (*See Belladonna*).

Liniment of Calamine (*See Zinc, Salts of*).

Liniment of Camphor (*See Camphor*).

Liniment of Camphor Ammoniated (*See Camphor and Ammonia*).

Liniment of Cantharides (*See Cantharides*).

Liniment of Chloroform (*See Chloroform*).

Liniment of Croton Oil (*See Croton Oil*).

Liniment of Mercury (*See Mercury, Ammonia and Camphor*).

Liniment of Opium (*See Opium*).

Liniment of Turpentine (*See Turpentine and Camphor*).

Mercury, Salts of.

Stomach Tube. Flour and Water; or Arrowroot and Water; or White of Egg and Water, *ad lib.* Leave in Stomach White of Egg and Water (1 egg = 4 grains of Corrosive Sublimate). Stimulants. Hot Bottles and Blankets.

Later. — Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically.

Morphine.

Stomach Tube or Emetic. Wash out Stomach at half hour intervals with Solution of Potassium Permanganate, diluted with five times the quantity of Warm Water, leaving about 5 ounces of the Diluted Solution in the Stomach. Mixture of Ammonia and Ether, 1 ounce; or Ether, 60 minims, hypodermically. Pinching. Flapping. Solution of Ammonia or Smelling Salts to nostrils. Faradism. Make patient walk about. Solution of Atropine Sulphate, 3 minims ($\frac{3}{100}$ grain of Atropine Sulphate), hypodermically, to be repeated in half an hour; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth or rectum. Capsule of Amyl Nitrite for inhalation. Artificial respiration.

Muriatic Acid (Hydrochloric Acid, Spirits of Salts).

Introduce Water and use Stomach Tube with care. Sodium Bicarbonate, 2 drachms in 5 ounces of Water; or Soap and Water; or Chalk or Calcined Magnesia, $\frac{1}{2}$ ounce in $\frac{1}{2}$ pint of Water or Milk; or Solution of Ammonia or Sal Volatile, freely diluted; or White of Egg and Water; or Milk.

Later.—Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically, for pain, and Ice for thirst.

Mussels.

Emetic. Stimulants. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically. Castor Oil, 2 ounces, with Tincture of Opium, 10 minims.

Nepenthe.

Stomach Tube or Emetic. Wash out Stomach at half hour intervals with Solution of Potassium Permanganate, diluted with five times the quantity of Warm Water, leaving about 5 ounces of the Diluted Solution in the Stomach. Mixture of Ammonia and Ether, 1 ounce; or Ether, 60 minims, hypodermically. Pinching. Flapping. Solution of Ammonia or Smelling Salts to Nostrils. Faradism. Make patient walk about. Solution of Atropine Sulphate, 3 minims ($\frac{3}{100}$ grain of Atropine Sulphate), hypodermically, to be repeated in half an hour; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth or rectum. Capsule of Amyl Nitrite for inhalation. Artificial respiration.

Nicotine.

Immediate Treatment.—Stomach Tube or Emetic. Stimulants. Warmth. Artificial respiration. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically.

Later.—Stimulants.

Nitric Acid.

Immediate Treatment.—Soap and Water. Calcined Magnesia, Chalk or Whiting, $\frac{1}{2}$ ounce in $\frac{1}{2}$ pint of Water or Milk; or Sodium Bicarbonate, 2 drachms in 5 ounces of Water.

Later.—Demulcents. Olive Oil. White of Egg and Water. Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically.

Nitro-Benzene (Nitro-Benzol, Artificial Oil of Bitter Almond, Essence of Mirbane).

Stomach Tube or Emetic. Stimulants. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically. Artificial respiration. Faradism.

Nitro-Glycerin.

Immediate Treatment. — Recumbent position. Cold Affusion.

Later.—Hypodermic injection of Ergot, 3 minims; or Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically.

Nitrous Oxide Gas.

Insert finger into mouth and pharynx to make sure that there is no obstruction, *e.g.*, artificial teeth, etc. Extend the head and push the jaw forward. Pull tongue out with forceps. Fresh air. Loosen clothes. Place in Recumbent position. Artificial respiration. Flapping with wet towels. Inhalation of Oxygen.

Opium.

Stomach Tube or Emetic. Wash out Stomach at half hour intervals with Solution of Potassium Permanganate, diluted with five times the quantity of Warm Water, leaving about 5 ounces of the Diluted Solution in the Stomach. Mixture of Ammonia and Ether, 1 ounce; or Ether, 60 minims, hypodermically. Pinching. Flapping. Solution of Ammonia or Smelling Salts to

nostrils. Faradism. Make patient walk about. Solution of Atropine Sulphate, 3 minims ($\frac{3}{100}$ grain of Atropine Sulphate), hypodermically, to be repeated in half an hour; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth or rectum. Capsule of Amyl Nitrite for inhalation. Artificial respiration.

Oxalic Acid.

Stomach Tube with care. Wash out Stomach with Lime Water. Chalk, $\frac{1}{2}$ ounce in 5 ounces of Water; or Saccharated Lime Water, $\frac{1}{2}$ ounce in 2 ounces of Water, repeated every half hour for 8 doses. Stimulants, hypodermically.

Paraffin Oil.

Emetics. Stimulants. Hot Bottles and Blankets. Friction.

Paraldehyde.

Immediate Treatment.—Prevent sleep. Stomach Tube and Emetic. Hot Blankets. Hot Bottle to feet.

Later.—Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Artificial respiration. Faradism. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically. Capsule of Amyl Nitrite for inhalation. Inhalation of Oxygen.

Paris Green (*See Arsenic*).

Petroleum.

Emetics. Stimulants. Friction. Hot Bottles and Blankets.

Phenol (Carbolic Acid).

Stomach Tube. Sodium Sulphate, 1 ounce in 3 ounces of Water; or Saccharated Lime Water, 1 ounce in 3 ounces of Water.

Later.—Demulcents. Olive Oil. Brandy, subcutaneously or by enema. Warmth.

Phosphorus.

Stomach Tube or Emetic. Solution of Hydrogen Peroxide, 30 minims, repeated. Purgative of $\frac{1}{2}$ ounce of Magnesium Sulphate in 4 ounces of Water. Avoid Oils, Fats and Milk.

Physostigmine.

Emetics. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water. Solution of Atropine Sulphate, 2 minims ($\frac{1}{50}$ grain of Atropine Sulphate), hypodermically; or Tincture of Belladonna, 15 minims in 1 ounce of Water, by mouth every 15 minutes until pupils dilate. Chloral Hydrate, 10 grains in 2 ounces of Water, by mouth or rectum every 15 minutes for 4 doses. Stimulants. Artificial respiration. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically.

Prussic Acid (*See Hydrocyanic Acid*).

Pyroligneous Acid (*See Acetic Acid*).

Rat Pastes (Arsenic).

Immediate Treatment.—Stomach Tube or Emetic. Ferric Hydrate (prepared by mixing Solution of Ferric Chloride, ½ ounce in half a tumbler of Water with an equal quantity of Saturated Solution of Washing Soda). Collect precipitate on a handkerchief and mix with half a tumbler of Hot Water. Repeat frequently. (Solution of Ammonia may be used instead of Washing Soda).

Note: Roth and Ringeisen's contains Arsenic and Phosphorus.

Rough on Rats and *Simpson's* contain Arsenic.

Rat Pastes (Phosphorus).

Stomach Tube or Emetic. Solution of Hydrogen Peroxide, 30 minims, repeated. Purgative of ½ ounce of Magnesium Sulphate in 4 ounces of Water. Avoid Oils, Fats and Milk.

Note: James's, Sanford's and Steiner's contain Phosphorus.

Roth and Ringeisen's contains Phosphorus and Arsenic.

Salts of Lemon.

Stomach Tube with care. Wash out Stomach with Lime Water. Chalk, ½ ounce in 5 ounces of Water; or Saccharated Lime Water, ½ ounce in 2 ounces of Water, repeated every half hour for 8 doses. Stimulants, hypodermically.

Salts of Sorrel.

Stomach Tube with care. Wash out Stomach with Lime Water. Chalk, ½ ounce in 5 ounces of Water; or Saccharated Lime Water, ½ ounce in 2 ounces of Water, repeated every half hour for 8 doses. Stimulants, hypodermically.

Santonin.

Immediate Treatment.—Stomach Tube or Emetic. Stimulants. If convulsions, either Chloral Hydrate, 15 grains in 2 ounces of Water; or Potassium Bromide, 30 grains in 2 ounces of Water.

Later.—Stimulants.

Savin.

Immediate Treatment.—Emetic.

Later.—Demulcents. Castor Oil. Morphine Sulphate, ¼ grain, hypodermically.

Scheele's Green (*See Arsenic*).

Sewer Gas.

Fresh air. Artificial respiration. Stimulants. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Hot Bottles and Blankets. Faradism.

Silver, Salts of.

Common Salt, ½ ounce in a pint of Water. Stomach Tube. White of Egg and Water. Milk.

Snake Bites.

Proximal Ligature. Remove ligature at intervals for a second or two, and re-apply, so as to allow only very small quantities of the poison to get into the circulation and yet avoid gangrene. Incision. Cauterisation. Inject into the puncture made by the fangs Solution of Potassium Permanganate, 40 minims; or Condy's Red Fluid, 20 minims; or 20 minims of a Solution consisting of 1 part of Solution of Potash and 6 parts of Water; or Solution of Ammonia, 10 minims; or 40 minims of a 2 % Solution of Calcium Hypochlorite. Stimulants very freely. Bleeding and Intravenous injection of Warm Saline Solution. Artificial respiration. Solution of Strychnine Hydrochloride, 5 minims ($\frac{1}{20}$ grain of Strychnine Hydrochloride), hypodermically, to be repeated until symptoms of Strychnine spasm appear. For the bite of a Rattle Snake, Olive Oil freely by mouth, and also to be rubbed into the skin.

Soldering Fluid (*See Zinc Chloride*).

Spirits of Salts (Hydrochloric Acid, Muriatic Acid).

Introduce Water and use Stomach Tube with care. Sodium Bicarbonate, 2 drachms in 5 ounces of Water; or Soap and Water; or Chalk or Calcined Magnesia, $\frac{1}{2}$ ounce in $\frac{1}{2}$ pint of Water or Milk; or Solution of Ammonia or Sal Volatile, freely diluted; or White of Egg and Water or Milk.

Later.—Morphine Sulphate, $\frac{1}{4}$ grain, hypodermically, for pain, and Ice for thirst.

Strychnine.

Immediate Treatment.—Wash out Stomach at half hour intervals with Solution of Potassium Permanganate, diluted with three times the quantity of Warm Water. Stomach Tube and Emetic. Injection of Apomorphine, 5 minims, hypodermically. Potassium Bromide, 1 drachm in 1 ounce of Water every half hour. Chloral Hydrate, 5 grains in 2 drachms of Water, hypodermically, or in 1 ounce of Water, by rectum, repeated.

Later.—Inhalation of Chloroform if convulsions are very severe. Artificial respiration.

Sulphuretted Hydrogen.

Fresh air. Artificial respiration. Stimulants. Strong Hot Coffee, 1 pint; or Caffeine Sodium Salicylate, 3 grains, hypodermically. Hot Bottles and Blankets. Faradism.

Sulphuric Acid.

Immediate Treatment.—Soap and Water. Calcined Magnesia, Chalk or Whiting, ½ ounce in ½ pint of Water or Milk; or Sodium Bicarbonate, 2 drachms in 5 ounces of Water.

Later.—Demulcents. Olive Oil. White of Egg and Water. Morphine Sulphate, ¼ grain, hypodermically.

Tartar Emetic.

Stomach Tube. Glycerin of Tannic Acid, 2 drachms in 5 ounces of Water; or Tannic Acid, 30 grains in 5 ounces of Water. Vegetable astringents. Strong Tea or Coffee, ½ pint.

Later.—Stimulants.

Turpentine.

Stomach Tube or Emetic. Magnesium Sulphate, ½ ounce in 2 ounces of Water. Demulcents.

Later.—Morphine Sulphate, ¼ grain, hypodermically.

Verdigris.

Immediate Treatment. — White of Egg and Water. Milk. Stomach Tube if vomiting has not occurred.

Later.—Demulcents. Morphine Sulphate, ¼ grain, hypodermically, and Hot Fomentations if much pain.

Vermin Killers (Arsenic).

Immediate Treatment.—Stomach Tube or Emetic. Ferric Hydrate (prepared by mixing Solution of Ferric Chloride, ½ ounce in half a tumbler of Water, with an equal quantity of Saturated Solution of Washing Soda). Collect precipitate on a handkerchief and mix with half a tumbler of Hot Water. Repeat frequently. (Solution of Ammonia may be used instead of Washing Soda).

Note : Roth and Ringeisen's contains Arsenic and Phosphorus.

Rough on Rats and *Simpson's* contain Arsenic.

Vermin Killers (Phosphorus).

Stomach Tube or Emetic. Solution of Hydrogen Peroxide, 30 minims, repeated. Purgative of ½ ounce of Magnesium Sulphate in 4 ounces of Water. Avoid Oils, Fats and Milk.

Note: Roth and Ringeisen's contains Phosphorus and Arsenic.

Vermin Killers (Strychnine).

Immediate Treatment.—Wash out Stomach at half hour intervals with Solution of Potassium Permanganate, diluted with three times the quantity of Warm Water. Stomach Tube and Emetic. Injection of Apomorphine, 5 minims, hypodermically. Potassium Bromide, 1 drachm in 1 ounce of Water every half hour. Chloral Hydrate, 5 grains in 2 drachms of Water, hypodermically, or in 1 ounce of Water by rectum, repeated.

Later.—Inhalation of Chloroform if convulsions are very severe. Artificial respiration.

Note: Battle's, Butler's and Gibson's contain Strychnine.

Vitriol, Blue (Copper Sulphate).

Immediate Treatment. — White of Egg and Water. Milk. Stomach Tube if vomiting has not occurred.

Later.—Demulcents. Morphine Sulphate, ¼ grain, hypodermically, and Hot Fomentations, if much pain.

Vitriol, White (Zinc Sulphate).

White of Egg and Water. Washing Soda in dilute solution. Demulcents. Strong Tea, ½ pint. Hot Fomentations. Morphine Sulphate, ¼ grain, hypodermically.

Weed Killers (*See Arsenic*).

White Precipitate.

Stomach Tube. Flour and Water; or Arrowroot and Water; or White of Egg and Water, *ad lib.* Leave in Stomach White of Egg and Water (1 egg = 4 grains of Corrosive Sublimate). Stimulants. Hot Bottles and Blankets.

Later.—Morphine Sulphate, ¼ grain, hypodermically.

Zinc, Salts of.

White of Egg and Water. Washing Soda in dilute solution. Demulcents. Strong Tea, ½ pint. Hot Fomentations. Morphine Sulphate, ¼ grain, hypodermically.

PHARMACOPŒIA.

—o—

BALNEA.

Balneum Calidum.

Water at 98°F (36°C) to 110°F (43°C).

Balneum Frigidum.

Water at 60°F (15°C).

The temperature may be reduced by the addition of Ice if necessary.

Balneum Tepidum.

Water at 85°F (29°C) to 92°F (33°C).

To make the following Medicated Baths, add the quantity of ingredient mentioned to 30 gallons of Water at 98°F (36°C):—

Balneum Alkalinum.

Sodium Carbonate	4, 6 or 8 ounces.

Balneum Alkalinum Compositum.

Borax	1½ ounces.
Sodium Carbonate	6 ounces.
Starch, in the form of paste ...	4 ounces.

Balneum Bituminis.

Solution of Coal Tar ...	8 fluid ounces.

Balneum Creosoti.

Creosote	2 fluid drachms.
Glycerin	2 fluid ounces.

Balneum Furfuris.

Wheaten Bran	4 pounds.

Tie the Bran in a muslin bag and macerate with frequent agitation for 15 minutes in one third of the Water at 180°F (82°C); then add more Water to produce the requisite quantity at 98°F (36°C).

Balneum Sinapis.

Mustard	2 pounds.

Rub the Mustard to a smooth paste with Cold Water before adding it to the Hot Water.

Balneum Sodii Chloridi.

Salt	8 pounds.

Balneum Sulphuratum.

Sulphurated Potash	4 ounces.

COLLODIUM.

Collodium Acidi Salicylici.

Salicylic Acid	1 drachm.
Extract of Indian Hemp	20 grains.
Flexible Collodion	1 fluid ounce.

COLLUNARIA.

The following Nasal Douches are to be diluted with two parts of Warm Water before use:—

Collunarium Alkalinum.

Borax	6 grains.
Sodium Bicarbonate...	6 grains.
Liquefied Carbolic Acid	2 minims.
Sugar	10 grains.
Water...	to 1 fluid ounce.

Collunarium Ammonii Chloridi.

Ammonium Chloride	10 grains.
Sodium Chloride	6 grains.
Sugar	10 grains.
Water...	to 1 fluid ounce.

Collunarium Boracis.

Borax	4 grains.
Sodium Bicarbonate...	4 grains.
Sodium Chloride	4 grains.
Sugar	10 grains.
Water...	to 1 fluid ounce.

Collunarium Potassii Permanganatis.

Potassium Permanganate	$\frac{1}{4}$ grain.
Sodium Chloride	10 grains.
Water	to 1 fluid ounce.

CONFECTIONES.

Confectio Guaiaci Composita.

Guaiacum Resin in powder ...	½ ounce.
Mustard	1 ounce.
Potassium Nitrate in powder ...	¼ ounce.
Rhubarb Root in powder ...	¼ ounce.
Sublimed Sulphur	1 ounce.
Treacle	to 16 ounces.

Dose :—1 drachm.

Confectio Sennæ et Piperis.

Confection of Senna ... } of each equal parts.
Confection of Pepper ... }

Dose :—1 to 2 drachms.

Confectio Sennæ et Sulphuris.

Confection of Senna ... } of each equal parts.
Confection of Sulphur ... }

Dose :—1 to 2 drachms.

DENTIFRICIA.

Dentifricium Acidi Carbolici Compositum.

Liquefied Carbolic Acid	15 minims.
Orris Root in powder	30 grains.
Refined Sugar in powder ...	30 grains.
Precipitated Calcium Carbonate...	1 ounce.

Dentifricium Eucalypti Compositum.

Oil of Eucalyptus	15 minims.
Powdered Castile Soap	1 drachm.
Precipitated Chalk	1 ounce.

Dentifricium Potassii Chloratis.

Powdered Potassium Chlorate	... 16 grains.
Compound Eucalyptus Dentifrice	... 4 drachms.
Glycerin	of each equal parts—
Water	a sufficiency.

EMPLASTRUM.

Emplastrum Acidi Salicylici et Creosoti.

This is of two strengths, and contains 10 or 50 grammes of Salicylic Acid to 20 or 50 grammes of Creosote respectively in $\frac{1}{5}$ square metre.

EMULSIONES.

Emulsio Chloroformi.

Chloroform	1 fluid ounce.
Tincture of Quillaia ...	1 fluid drachm.
Water...	to 1 pint.

(To be used instead of Spirit of Chloroform).

Emulsio Iodoformi.

Iodoform in powder	1 ounce.
Glycerin	...to 10 fluid ounces.

ENEMATA

Enema Amyli.

Powdered Starch	... 2 drachms.
Water	5 fluid ounces.

Rub the Starch to a smooth paste with the Water and heat the Mixture to boiling.

Enema Asafetidæ.

Tincture of Asafetida ...	½ fluid ounce.
Starch Enema	to 4 fluid ounces.

Enema Chloral.

Chloral Hydrate	10 to 40 grains.
Starch Enema	to 4 fluid ounces.

Enema Glycerini.

1 to 4 fluid drachms of Glycerin to be injected with or without an equal quantity of Warm Water.

Enema Nutriens.

The basis of the Enema should consist of Peptonised Milk prepared as on *page* 85. To this may be added 2 ounces of Peptone (Somatose), Grape Sugar or Raw Starch; or 3 Eggs may be beaten up with the Milk. If Eggs be used, 15 grains of Salt should be added for each Egg, as this is found to greatly promote absorption. Alcohol may also be added, preferably in the form of Red Wine (2 to 4 fluid ounces). The total bulk of the Enema should not exceed 9 fluid ounces. It should be introduced at the body temperature through a small bore Œsophageal Tube, passed as high up as possible and connected with a funnel raised to a height not exceeding 3 feet. If badly retained, add 15 to 20 minims of Tincture of Opium. Not more than three such Enemata should be given in 24 hours.

Enema Olei Ricini.

Castor Oil	1 fluid ounce.
Soft Soap	½ ounce.
Warm Water	10 fluid ounces.

Enema Oleosum.

Olive Oil	4 fluid ounces.
Soft Soap	½ ounce.
Warm Water	16 fluid ounces.

Olive Oil may be given alone, from ½ to 1 pint. warmed to a temperature of 90°F (32°C).

Enema Opii.

Tincture of Opium	10 to 60 minims.
Starch Enema	to 4 fluid ounces.

Enema Simplex.

Soft Soap	1 ounce.
Warm Water	to 1 pint.

Enema Terebinthinæ.

Turpentine	½ to 2 fluid drachms.
Starch Enema	to 1 pint.

EXTRACTA.

Extractum Malti.

Dose :—1 to 4 fluid drachms.

Extractum Malti cum Oleo Morrhuæ.

Extract of Malt with about 20% of Cod Liver Oil.

Dose :—1 to 4 fluid drachms.

FOTUS.

Fotus Acidi Borici.

Dip Boric Lint in Hot Water, lightly wring out and apply to the affected part. Cover with Gutta Percha Tissue, Cotton Wool and Bandage.

Fotus Belladonnæ.

Paint the affected part with Glycerin of Belladonna, and apply Flannel wrung out of Hot Water. Cover with Gutta Percha Tissue, Cotton Wool and Bandage.

Fotus Opii.

Sprinkle Tincture of Opium on Flannel wrung out of Hot Water, and apply to the affected part. Cover with Gutta Percha Tissue, Cotton Wool and Bandage.

Fotus Terebinthinæ.

Sprinkle Turpentine on Flannel wrung out of Hot Water, and apply to the affected part. Cover with Gutta Percha Tissue, Cotton Wool and Bandage.

GARGARISMATA.

Gargarisma Acidi Carbolici.

Glycerin of Carbolic Acid ...	25 minims.
Water	to 1 fluid ounce.

Gargarisma Acidi Tannici.

Glycerin of Tannic Acid...	1 fluid drachm.
Water	to 1 fluid ounce.

Gargarisma Aluminis.

Alum	10 grains.
Water	to 1 fluid ounce.

Gargarisma Aluminis Compositum.

Alum	5 grains.
Diluted Sulphuric Acid	10 minims.
Tincture of Myrrh	10 minims.
Water	to 1 fluid ounce.

Gargarisma Boracis.

Borax	10 grains.
Tincture of Myrrh	10 minims.
Water	to 1 fluid ounce.

Gargarisma Chlori.

Potassium Chlorate	10 grains.
Strong Hydrochloric Acid ...	3 minims.
Water	to 1 fluid ounce.

Place the Potassium Chlorate in a *dry* bottle, pour the Acid upon it, cork loosely, and set aside for 10 minutes. Then add the Water in 3 or 4 successive portions, shaking between each addition.

Gargarisma Potassii Chloratis.

Potassium Chlorate	10 grains.
Water	to 1 fluid ounce.

Gargarisma Potassii Chloratis Compositum

Potassium Bicarbonate	5 grains.
Sodium Bicarbonate	5 grains.
Potassium Chlorate Gargle...	to 1 fluid ounce.

Gargarisma Potassii Chloratis cum Ferro.

Solution of Ferric Chloride ...	15 minims.
Potassium Chlorate Gargle	to 1 fluid ounce.

Gargarisma Potassii Permanganatis.

Solution of Potassium Permanganate	10 minims.
Water	to 1 fluid ounce.

Gargarisma Sodæ Chlorinatæ.

Solution of Chlorinated Soda ...	40 minims.
Water	to 1 fluid ounce.

GLYCERINA.

Glycerinum Belladonnæ.

Alcoholic Extract of Belladonna...	2 drachms.
Water	1 fluid drachm.
Glycerin	1 fluid ounce.

Glycerinum Ferri Perchloridi.

Solution of Ferric Chloride	1 fluid drachm.
Glycerin	to 1 fluid ounce.

Glycerinum Hydrargyri Perchloridi (1 in 1,000).

Mercuric Chloride...	17½ grains.
Glycerin	1 pint.
Distilled Water	to 2 pints.

GUTTÆ.

GUTTÆ ACIDI CARBOLICI.

(*Aural*).

Glycerin of Carbolic Acid ...	of each
Glycerin	equal parts.

GUTTÆ ATROPINÆ SULPHATIS.

(*Ophthalmic*).

Atropine Sulphate	...	1, 2 or 4 grains.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ COCAINÆ HYDROCHLORIDI.

(*Ophthalmic*).

Cocaine Hydrochloride ...	...	10 grains.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ CUPRI SULPHATIS.

(*Ophthalmic*).

Copper Sulphate ...	...	... 2 grains.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ DUBOISINÆ SULPHATIS.

(*Ophthalmic*).

Duboisine Sulphate	...	... 1 grain.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ ESERINÆ SULPHATIS.

(*Ophthalmic*).

Eserine Sulphate ...	...	½, 1, 2 or 4 grains.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ ESERINÆ ET COCAINÆ.

(*Ophthalmic*).

Eserine Sulphate ...	...	... 1 grain.
Cocaine Hydrochloride ...	...	4 grains.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ HOMATROPINÆ HYDROBROMIDI.

(*Ophthalmic*).

Homatropine Hydrobromide	...	4 grains.
Distilled Water		to 1 fluid ounce.

GUTTÆ HOMATROPINÆ ET COCAINÆ.

(*Ophthalmic*).

Homatropine Hydrobromide	...	6 grains.
Cocaine Hydrochloride ...	...	10 grains.
Distilled Water ...	...	to 1 fluid ounce.

GUTTÆ HOMATROPINÆ ET COCAINÆ OLEOSÆ.

(*Ophthalmic*).

Homatropine ...	...	10 grains.
Cocaine	...	10 grains.
Castor Oil	...	to 1 fluid ounce.

GUTTÆ HYDRARGYRI NITRATIS.

(*Aural*).

Diluted Nitrate of Mercury Ointment		1 drachm.
Olive Oil	...	½ fluid ounce.
Liquid Paraffin		to 1 fluid ounce.

GUTTÆ HYDROGENII PEROXIDI.

(*Aural*).

Solution of Hydrogen Peroxide (10 vols.)	of each equal parts.
Distilled Water	

GUTTÆ HYOSCINÆ HYDROBROMIDI.

(*Ophthalmic*).

Syn: **Guttæ Scopolaminæ.**

Hyoscine Hydrobromide ...		1 or 2 grains.
Distilled Water		to 1 fluid ounce.

GUTTÆ PILOCARPINÆ NITRATIS.

(*Ophthalmic*).

Pilocarpine Nitrate ...	... 2 or 4 grains.
Distilled Water	to 1 fluid ounce.

GUTTÆ ZINCI CHLORIDI.

(*Ophthalmic*).

Zinc Chloride	1 grain.
Distilled Water	to 1 fluid ounce.

GUTTÆ ZINCI CHLORIDI ET ACIDI BORICI.

(*Ophthalmic*).

Zinc Chloride	½, 1 or 2 grains.
Boric Acid...	10 grains.
Distilled Water	to 1 fluid ounce.

GUTTÆ ZINCI CHLORIDI ET COCAINÆ.

(*Ophthalmic*).

Zinc Chloride	2 grains.
Cocaine Hydrochloride	10 grains.
Distilled Water	to 1 fluid ounce.

HAUSTUS.

Haustus Aperiens.

Syn: **H.D.**

Magnesium Sulphate	150 grains.
Emulsion of Chloroform ...	7½ minims.
Infusion of Senna... ...	to 1 fluid ounce.

Haustus Bromo-Chloral Compositus.

Potassium Bromide	10 grains.
Chloral Hydrate	10 grains.
Juice of Hyoscyamus	10 minims.
Liquid Extract of Liquorice ...	2 minims.
Syrup	10 minims.
Tincture of Indian Hemp ...	3 minims.
Tincture of Orange	3 minims.
Water	to 1 fluid ounce.

Haustus Filicis.

Liquid Extract of Male Fern ...	1 fluid drachm.
Syrup of Ginger	1 fluid drachm.
Tincture of Quillaia	30 minims.
Peppermint Water ...	to 1 fluid ounce.

INJECTIONES

The following Hypodermic Discs are kept in stock:—

Apomorphinæ Hydrochloridum ...	$\frac{1}{15}$, $\frac{1}{10}$ grain.
Atropinæ Sulphas ...	$\frac{1}{150}$, $\frac{1}{100}$, $\frac{1}{60}$ grain.
Caffeinæ Sodio-Salicylas ...	$\frac{1}{2}$, 1, 2 and 3 grains.
Digitalinum	$\frac{1}{100}$ grain.
Ergotinæ Citras	$\frac{1}{200}$, $\frac{1}{100}$ grain.
Eucainæ Hydrochloridum ...	$\frac{1}{3}$, 1 grain.
Hyoscinæ Hydrobromidum	$\frac{1}{200}$, $\frac{1}{100}$ grain.
Morphinæ Sulphas ...	$\frac{1}{12}$, $\frac{1}{8}$, $\frac{1}{6}$, $\frac{1}{4}$, $\frac{1}{3}$, $\frac{1}{2}$ grain.
Pilocarpinæ Nitras ...	$\frac{1}{10}$, $\frac{1}{6}$, $\frac{1}{3}$, $\frac{1}{2}$ grain.
Strychninæ Sulphas ...	$\frac{1}{100}$, $\frac{1}{60}$, $\frac{1}{30}$ grain.

Injectio Acidi Borici.

(*Vaginal*).

Boric Acid...	1 drachm.
Water	1 pint.

Injectio Acidi Tannici.

(*Vaginal*).

Tannic Acid	1 drachm.
Water	1 pint.

Injectio Aluminis.

(*Vaginal*).

Alum	1 drachm.
Water	1 pint.

Injectio Aluminis Composita.

(*Urethral*).

Alum	4 grains.
Carbolic Acid	$\frac{1}{4}$ grain.
Water	1 fluid ounce.

Injectio Plumbi.

(*Vaginal*).

Strong Solution of Lead Sub-acetate	1 fluid drachm.
Water	1 pint.

Injectio Potassii Permanganatis.

(*Vaginal*).

Potassium Permanganate ...	5 grains.
Water	1 pint.

Injectio Protargolis.

(*Urethral*).

Protargol ... ... $\frac{1}{2}$ to 1 % dissolved in water.

Injectio Zinci Chloridi.

(*Urethral*).

Zinc Chloride	1 grain.
Water	1 fluid ounce.

Injectio Zinci Chloridi.

(*Vaginal*).

Zinc Chloride	5 grains.
Water	1 pint.

Injectio Zinci Permanganatis.

(*Urethral*).

Zinc Permanganate	½ grain.
Water	1 fluid ounce.

Injectio Zinci Sulphatis.

(*Urethral*).

Zinc Sulphate	2 grains.
Water	1 fluid ounce.

Injectio Zinci Sulphatis.

(*Vaginal*).

Zinc Sulphate	1 drachm.
Water	1 pint.

Injectio Zinci Sulphocarbolatis.

(*Urethral*).

Zinc Sulphocarbolate	5 grains.
Water	1 fluid ounce.

LINCTUS.

Linctus Camphoræ Compositus.

Compound Tincture of Camphor ...	20 minims.
Syrup of Squills	20 minims.
Syrup of Tolu	20 minims.

Linctus Morphinæ Compositus.

Solution of Morphine Hydrochloride	6 minims.
Cherry Laurel Water	8 minims.
Spirit of Chloroform	4 minims.
Syrup of Lemons	30 minims.
Glycerin	to 1 fluid drachm.

Linctus Scillæ.

Syrup of Squills	15 minims.
Syrup of Lemons...	15 minims.
Syrup of Poppies	15 minims.
Syrup of Tolu	15 minims.

Linctus Terp=Heroin Compositus.

Terpene Hydrate	$\frac{1}{8}$ grain.
Heroin Hydrochloride	$\frac{1}{24}$ grain.
Menthol	$\frac{1}{24}$ grain.
Alcohol 90%	a sufficiency.
Glycerin	20 minims.
Syrup of Lemons	10 minims.
Syrup of Tolu	12$\frac{1}{2}$ minims.
Syrup	to 1 fluid drachm.

LINIMENTA.

Linimentum A. B. C.

Liniment of Aconite ...
Liniment of Belladonna } of each equal parts.
Liniment of Chloroform

Linimentum Aconiti cum Sapone.

Liniment of Aconite	1 part.
Liniment of Soap	3 parts.

Linimentum Belladonnæ cum Sapone.

Liniment of Belladonna	1 part.
Liniment of Soap	3 parts.

Linimentum Calaminæ.

Calamine	35 grains.
Solution of Lime ...	½ fluid ounce.
Olive Oil	to 1 fluid ounce.

Linimentum Cantharidis.

Blistering Liquid	1 fluid drachm.
Glacial Acetic Acid ...	1 fluid drachm.
Methylated Spirit ...	1 fluid ounce.

Linimentum Mentholis.

Menthol	1 drachm.
Olive Oil	to 1 fluid ounce.

Linimentum Myristicæ.

Oil of Nutmeg	1 part.
Olive Oil	3 parts.

LIQUORES.

LIQUOR ARGENTI NITRATIS.

(*Ophthalmic*).

Silver Nitrate	5 or 10 grains.
Distilled Water	to 1 fluid ounce.

LIQUOR FLUORESCEIN.

(*Ophthalmic*).

Fluorescein	8 grains.
Sodium Bicarbonate...	12 grains.
Distilled Water	to 1 fluid ounce.

LIQUOR PROTARGOLIS.

(*Ophthalmic*).

Protargol	40, 60, 80 or 100 grains.
Distilled Water	to 1 fluid ounce.

Liquor Rubri.

Fuchsine	1 grain.
Methylated Spirit	½ fluid ounce.
Water...	to 1 fluid ounce.

LOTIONES.

LOTIO ACIDI BORICI.

(*Ophthalmic*).

Boric Acid	15 grains.
Distilled Water	to 1 fluid ounce.

Lotio Acidi Borici.

Boric Acid	15 grains.
Water...	to 1 fluid ounce.

Lotio Acidi Carbolici.

Carbolic Acid (Crystals)	1 ounce.
Water...	to 1 pint.

LOTIO ACIDI HYDROCYANICI CUM SODIO.

(*Ophthalmic*).

Diluted Hydrocyanic Acid ...	... 4 minims.
Borax	... 4 grains.
Sodium Bicarbonate ...	... 4 grains.
Distilled Water ...	... to 1 fluid ounce.

Lotio Acidi Sulphurosi.

Sulphurous Acid ...	... 2 fluid drachms.
Water	... to 1 fluid ounce.

LOTIO ALUMINIS.

(*Ophthalmic*).

Alum	... ... 4 grains.
Distilled Water ...	... to 1 fluid ounce.

LOTIO ALUMINIS COMPOSITA.

(*Ophthalmic*).

Alum	... 3 grains.
Zinc Sulphate	... 1 grain.
Distilled Water ...	... to 1 fluid ounce.

LOTIO BORACIS.

(*Ophthalmic*).

Borax	... ... 10 grains.
Sodium Bicarbonate...	... 10 grains.
Distilled Water ...	... to 1 fluid ounce.

Lotio Calaminæ.

Calamine	... 30 grains.
Zinc Oxide	... ... 20 grains.
Glycerin	... ... 30 minims.
Solution of Lime ...	... 0 1 fluid ounce.

Lotio Evaporans.

Ammonium Chloride	10 grains.
Methylated Spirit	2 fluid drachms.
Solution of Ammonium Acetate	1 fluid drachm.
Water	to 1 fluid ounce.

Lotio Glycerini Plumbi Subacetatis.

Glycerin of Lead Subacetate	... 30 minims.
Glycerin	... 30 minims.
Water	to 1 fluid ounce.

Lotio Hydrargyri Biniodidi.

(1 in 500).

Mercuric Iodide	... 17½ grains.
Potassium Iodide	... 12 grains.
Water	... to 1 pint.

LOTIO HYDRARGYRI PERCHLORIDI.

(*Ophthalmic*).

Mercuric Chloride	... 2 grains.
Distilled Water	... to 1 pint.

Lotio Hydrargyri Perchloridi.

(1 in 500).

Mercuric Chloride	... 17½ grains.
Water	... to 1 pint.

Lotio Picis Carbonis.

Solution of Coal Tar ...	... 10 minims.
Water	to 1 fluid ounce.

Lotio Plumbi Evaporans

Strong Solution of Lead Sub-acetate	... 6 minims.
Methylated Spirit	2 fluid drachms.
Water	to 1 fluid ounce

Lotio Plumbi cum Opio.

Extract of Opium	4 grains.
Diluted Solution of Lead Sub-acetate	to 1 fluid ounce.

Lotio Plumbi et Picis.

Strong Solution of Lead Sub-acetate	... 10 minims.
Solution of Coal Tar ...	... 20 minims.
Water...	to 1 fluid ounce.

LOTIO QUININÆ SULPHATIS.

(*Ophthalmic*).

Quinine Sulphate	4 grains.
Diluted Sulphuric Acid ...	a sufficiency.
Distilled Water	to 1 fluid ounce.

Lotio Resorcini.

Resorcin	20 grains.
Glycerin	20 minims.
Water	to 1 fluid ounce.

Lotio Saponis Alcoholica.

Soft Soap	1 ounce.
Methylated Spirit	1 fluid ounce.

Lotio Sodæ Chlorinatæ.

Solution of Chlorinated Soda	1 fluid drachm.
Water...	to 1 fluid ounce.

LOTIO SODII BICARBONATIS.

(*Ophthalmic*).

Sodium Bicarbonate... ...	... 15 grains.
Distilled Water	to 1 fluid ounce.

Lotio Sodii Hyposulphitis.

Sodium Hyposulphite	1 drachm.
Water	to 1 fluid ounce.

LOTIO ZINCI SULPHATIS.

(*Ophthalmic*).

Zinc Sulphate	1 or 2 grains.
Distilled Water	to 1 fluid ounce.

Lotio Zinci Sulphatis.

Zinc Sulphate	2 grains.
Water	to 1 fluid ounce.

Lotio Zinci Sulphidi.

Zinc Sulphide	12 grains.
Sulphurated Potash	12 grains.
Water	to 1 fluid ounce.

MISTURÆ.

Mistura Acidi Sulphurici cum Opio.

Aromatic Sulphuric Acid	15 minims.
Tincture of Opium	5 minims.
Compound Tincture of Cardamoms	20 minims.
Chloroform Water	to ½ fluid ounce.

Mistura Aloes Composita pro Infantibus.

Syn: **Anti-Constipation Mixture for Children.**

Tincture of Aloes	4 minims.
Tincture of Ginger	2 minims.
Tincture of Hyoscyamus	5 minims.
Tincture of Nux Vomica	½ minim.
Syrup of Senna	15 minims.
Water	to 1 fluid drachm.

Mistura Ammoniæ cum Æthere.

Aromatic Spirit of Ammonia	... 30 minims.
Spirit of Ether	... 30 minims.
Chloroform Water	to ½ fluid ounce

Mistura Ammoniæ et Ipecacuanhæ.

Ammonium Carbonate	3 grains.
Ipecacuanha Wine	... 20 minims.
Camphor Water	to ½ fluid ounce.

Mistura Ammoniæ et Ipecacuanhæ pro Infantibus.

Ammonium Carbonate	1 grain.
Ipecacuanha Wine	2½ minims.
Glycerin	... 10 minims.
Water...	... to 1 fluid drachm.

Mistura Aperiens pro Infantibus.

Magnesium Sulphate	3 grains.
Compound Tincture of Rhubarb	10 minims.
Caraway Water ...	... to 1 fluid drachm

Mistura Asafetidæ Composita.

Tincture of Asafetida	5 minims.
Ammonium Carbonate	4 grains.
Liquid Extract of Cascara Sagrada	10 minims.
Infusion of Valerian (1 in 40)	to ½ fluid ounce.

Mistura Bismuthi pro Infantibus.

Bismuth Oxycarbonate	5 grains.
Sodium Bicarbonate...	2 grains.
Caraway Water ...	... to 1 fluid drachm.

Mistura Bismuthi Astringens pro Infantibus.

Bismuth Oxycarbonate	5 grains.
Aromatic Powder of Chalk with Opium	1 grain.
Glycerin of Tannic Acid	5 minims.
Dill Water	to 1 fluid drachm.

Mistura Bismuthi cum Magnesia.

Bismuth Oxycarbonate	15 grains.
Light Magnesium Carbonate ...	20 grains.
Emulsion of Chloroform	10 minims.
Water	to ½ fluid ounce.

Mistura Bismuthi cum Opio.

Bismuth Oxycarbonate	15 grains.
Tincture of Opium	5 minims.
Peppermint Water	to ½ fluid ounce.

Mistura Bismuthi cum Soda.

Bismuth Oxycarbonate	15 grains.
Sodium Bicarbonate...	15 grains.
Emulsion of Chloroform	10 minims.
Water	to ½ fluid ounce.

Mistura Buchu Composita.

Potassium Citrate	20 grains.
Emulsion of Chloroform	10 minims.
Tincture of Hyoscyamus	20 minims.
Infusion of Buchu	to ½ fluid ounce.

Mistura Camphoræ cum Æthere.

Spirit of Ether	30 minims.
Camphor Water	to ½ fluid ounce

Mistura Capsici Sedativa.

Syn: **Anti-Alcoholic Mixture.**

Tincture of Capsicum ...	... 5 minims.
Tincture of Ginger ...	... ... 20 minims.
Potassium Bromide ...	... ... 10 grains.
Sodium Bicarbonate...	... ... 10 grains.
Decoction of Cinchona	... to ½ fluid ounce.

Mistura Capsici Sedativa cum Magnesii Sulphate.

Syn: **Aperient Anti-Alcoholic Mixture.**

Magnesium Sulphate ...	... 1 drachm.
Sedative Capsicum Mixture...	to ½ fluid ounce.

Mistura Carminativa pro Infantibus.

Aromatic Spirit of Ammonia	... 2 minims.
Glycerin	... 5 minims.
Sodium Bicarbonate...	... ... 3 grains.
Dill Water	... to 1 fluid drachm.

Mistura Cascaræ.

Liquid Extract of Cascara Sagrada	1 fluid drachm.
Liquid Extract of Liquorice	... 30 minims.
Aromatic Spirit of Ammonia	... 40 minims.
Chloroform Water	to ½ fluid ounce.

Mistura Cascaræ Aperiens.

Magnesium Sulphate ...	... 30 grains.
Cascara Sagrada Mixture ...	to ½ fluid ounce.

Mistura Cascaræ Composita.

Magnesium Sulphate ... ... 1 drachm.
Glycerin .. ..
Liquid Extract of Cascara Sagrada ... ...
Liquid Extract of Liquorice } of each 1 fluid drachm.
Tincture of Hyoscyamus ... ... 20 minims.
Tincture of Nux Vomica ... ... 5 minims.
Compound Decoction of Aloes to 1 fluid ounce.

Mistura Cinchonæ Acida.

Diluted Hydrochloric Acid ... ... 10 minims.
Decoction of Cinchona ... to ½ fluid ounce.

Mistura Colchici.

Colchicum Wine ... ... 15 minims.
Potassium Citrate ... ... 20 grains.
Peppermint Water ... ... to ½ fluid ounce.

Mistura Colchici cum Magnesii Sulphate.

Magnesium Sulphate ... ... 30 grains.
Colchicum Mixture ... ... to ½ fluid ounce.

Mistura Diaphoretica.

Solution of Ammonium Acetate 2 fluid drachms.
Spirit of Nitrous Ether ... ... 30 minims.
Camphor Water ... ... to ½ fluid ounce.

Mistura Ergotæ.

Liquid Extract of Ergot ... ... 30 minims.
Syrup of Ginger ... ... 30 minims.
Infusion of Orange Peel ... to ½ fluid ounce.

Mistura Ferri Perchloridi.

Solution of Ferric Chloride... ... 10 minims.
Syrup ... ... 30 minims.
Infusion of Quassia ... ... to ½ fluid ounce.

Mistura Ferri et Aloes.

Compound Decoction of Aloes	2 fluid drachms.
Compound Mixture of Iron	to 1 fluid ounce.

Mistura Ferri et Ammonii Acetatis.

Solution of Ferric Chloride...	... 15 minims.
Solution of Ammonium Acetate	2 fluid drachms.
Diluted Acetic Acid	... 30 minims.
Syrup	... 30 minims.
Water...	to ½ fluid ounce.

Mistura Ferri et Arsenici cum Quinina.

Hydrochloric Solution of Arsenic	3 minims.
Iron and Quinine Mixture ...	to ½ fluid ounce.

Mistura Ferri cum Magnesii Sulphate.

Ferrous Sulphate	3 grains.
Magnesium Sulphate	30 grains.
Diluted Sulphuric Acid	10 minims.
Peppermint Water	to ½ fluid ounce.

Mistura Ferri et Quininæ.

Solution of Ferric Chloride	... 10 minims.
Quinine Sulphate	1 grain.
Water...	to ½ fluid ounce.

Mistura Ferri et Strychninæ.

Solution of Ferric Chloride...	... 10 minims.
Solution of Strychnine Hydrochloride	5 minims.
Syrup	... 30 minims.
Water...	to ½ fluid ounce.

Mistura Ferri et Strychninæ cum Magnesii Sulphate.

Magnesium Sulphate ...	... 30 grains.
Iron and Strychnine Mixture	to ½ fluid ounce.

Mistura Gentianæ Alkalina.

Sodium Bicarbonate... ...	... 15 grains.
Emulsion of Chloroform ...	... 10 minims.
Infusion of Gentian	to ½ fluid ounce.

Mistura Gentianæ cum Rheo.

Gentian Root	... 5 grains.
Rhubarb Root	... 5 grains.
Ginger...	... 1 grain.
Sodium Bicarbonate... ...	... 10 grains.
Peppermint Water	to ½ fluid ounce.

Mistura Hydrargyri Iodidi.

Solution of Mercuric Chloride	1 fluid drachm.
Potassium Iodide	... 5 grains.
Peppermint Water ...	to ½ fluid ounce.

Mistura Hyoscyami cum Camphora.

Tincture of Hyoscyamus ...	... 30 minims.
Spirit of Camphor	... 15 minims.
Mucilage of Gum Acacia ...	... 30 minims.
Potassium Bromide	... 10 grains.
Water...	to ½ fluid ounce.

Mistura Lobeliæ et Potassii Iodidi.

Ethereal Tincture of Lobelia	... 15 minims.
Potassium Iodide	... 4 grains.
Tincture of Belladonna ...	... 5 minims.
Camphor Water	to ½ fluid ounce.

Mistura Magnesii Sulphatis Composita.

Magnesium Sulphate ...	... 1 drachm.
Light Magnesium Carbonate	... 15 grains.
Tincture of Ginger	... 20 minims.
Peppermint Water ...	... to ½ fluid ounce.

Mistura Olei Morrhuæ pro Infantibus.

Cod Liver Oil	½ fluid ounce.
Essential Oil of Almonds ...	... ½ minim.
Gluside	... $\frac{1}{20}$ grain.
Decoction of Irish Moss	... to 1 fluid ounce.

Dose:—1 to 4 fluid drachms or more.

Mistura Olei Ricini cum Opio.

Syn: **Anodyne Anti-Colic Mixture.**

Castor Oil	½ fluid ounce.
Tincture of Opium	... 10 minims.
Mucilage of Gum Acacia	... 1½ fluid drachms.
Cinnamon Water ...	... to 1 fluid ounce.

Mistura Olei Santali.

Oil of Sandal Wood ...	... 15 minims.
Mucilage of Gum Acacia ...	... 30 minims.
Peppermint Water ...	... to ½ fluid ounce.

Mistura Potassii Bicarbonatis pro Infantibus.

Potassium Bicarbonate ...	... 5 grains.
Potassium Citrate	... 5 grains.
Syrup of Orange	... 5 minims.
Tincture of Nux Vomica ...	... 2½ minims.
Infusion of Gentian ...	... to 1 fluid drachm.

Mistura Potassii Bicarbonatis cum Calumba.

Potassium Bicarbonate ...	... 20 grains.
Aromatic Spirit of Ammonia	... 20 minims.
Emulsion of Chloroform ...	... 10 minims.
Infusion of Calumba ...	to ½ fluid ounce.

Mistura Potassii Bromidi Composita.

Syn: **Anti-Climacteric Mixture.**

Potassium Bromide	... 10 grains.
Aromatic Spirit of Ammonia	... 20 minims.
Tincture of Nux Vomica ...	... 5 minims.
Infusion of Gentian	to ½ fluid ounce.

Mistura Potassii Citratis Effervescens.

Alkaline Solution.

Potassium Bicarbonate ...	... 20 grains.
Water...	to 1 fluid ounce.

Acid Solution.

Citric Acid	... 15 grains.
Water...	to ½ fluid ounce.

Mix the two solutions and drink during effervescence.

Mistura Potassii Iodidi et Ammonii Carbonatis.

Potassium Iodide	... 5 grains.
Ammonium Carbonate ...	... 4 grains.
Decoction of Cinchona ...	to ½ fluid ounce.

Mistura Quininæ cum Acido Hydrobromico.

Quinine Sulphate	... 1 grain.
Diluted Hydrobromic Acid...	... 30 minims.
Water...	to ½ fluid ounce.

Mistura Rhei cum Magnesia pro Infantibus.

Compound Tincture of Rhubarb	15 minims.
Oil of Caraway	$\frac{1}{32}$ minim.
Syrup of Ginger	15 minims.
Solution of Magnesium Carbonate	to 1 fluid drachm.

Mistura Rubra.

Red Solution...	1 minim.
Water...	to 1 fluid ounce.

Mistura Salina.

Potassium Nitrate	10 grains.
Solution of Ammonium Acetate	3 fluid drachms.
Spirit of Nitrous Ether ...	30 minims.
Water...	to $\frac{1}{2}$ fluid ounce.

Mistura Salina Antimonialis.

Solution of Ammonium Acetate	2 fluid drachms.
Spirit of Nitrous Ether ...	30 minims.
Antimonial Wine	20 minims.
Water...	to $\frac{1}{2}$ fluid ounce.

Mistura Salina cum Magnesii Sulphate.

Solution of Ammonium Acetate	2 fluid drachms.
Magnesium Sulphate ...	20 grains.
Water...	to $\frac{1}{2}$ fluid ounce.

Mistura Scillæ Composita.

Vinegar of Squill	15 minims.
Compound Tincture of Camphor	20 minims.
Emulsion of Chloroform ...	10 minims.
Infusion of Gentian	to $\frac{1}{2}$ fluid ounce.

Mistura Scoparii Composita.

Potassium Acetate	... 20 grains.
Spirit of Juniper	... 20 minims.
Spirit of Nitrous Ether ...	... 30 minims.
Infusion of Broom	to ½ fluid ounce.

Mistura Senegæ Composita.

Ammonium Carbonate	4 grains.
Tincture of Squill	... 5 minims.
Infusion of Senega	to ½ fluid ounce.

Mistura Sodii Phosphatis Acidi.

Sodium-Dihydrogen Phosphate	... 30 grains.
Red Mixture	to ½ fluid ounce.

Mistura Sodii Phosphatis Acidi Composita.

Hexamethylenetetramine	5 grains.
Sodium Acid Phosphate Mixture to	½ fluid ounce.

Mistura Sodii Salicylatis et Potassii Citratis.

Sodium Salicylate	... 10 grains.
Potassium Citrate	... 20 grains.
Saturated Aqueous Solution of Menthol	to ½ fluid ounce.

NEBULÆ.

NEBULA ACIDI CARBOLICI.

(Throat).

Liquefied Carbolic Acid ...	... 3 minims.
Borax	... 15 grains.
Sodium Bicarbonate	... 15 grains.
Glycerin	... 40 minims.
Water	to 1 fluid ounce.

NEBULA ACIDI TANNICI.

(Throat).

Tannic Acid	5 grains.
Glycerin	1 fluid drachm.
Water	to 1 fluid ounce.

Nebula Cocainæ cum Menthole.

Cocaine	5 grains.
Menthol	10 grains.
Liquid Paraffin	to 1 fluid ounce.

Nebula Eucalypti.

Oil of Eucalyptus	... 10 minims.
Liquid Paraffin	to 1 fluid ounce.

NEBULA HYDRARGYRI PERCHLORIDI.

(Throat).

Cocaine Hydrochloride ...	... 8 grains.
Glycerin	1 fluid drachm.
Solution of Mercuric Chloride (1 in 1.000)	to 1 fluid ounce.

Nebula Mentholis.

Menthol	... 10 grains.
Liquid Paraffin	1 fluid ounce.

NEBULA POTASSII CHLORATIS CUM FERRO

(Throat).

Potassium Chlorate	... 15 grains.
Solution of Ferric Chloride...	... 20 minims.
Glycerin	1 fluid drachm.
Water	to 1 fluid ounce.

PASTÆ.

Pasta Resorcini.

Resorcin	20 grains.
Starch	22 grains.
Zinc Oxide	22 grains.
Hydrous Wool Fat ... } Soft Paraffin ... ... }	of each equal parts to 1 ounce.

Pasta Zinci.

Zinc Oxide	24 parts.
Salicylic Acid in powder	2 parts.
Starch...	24 parts.
Soft Paraffin	50 parts.

Pasta Zinci et Gelatini.

Syn: **Unna's Paste.**

Zinc Oxide	1 ounce.
Gelatine	1 ounce.
Water...	2½ fluid ounces.
Glycerin	2½ fluid ounces.

PIGMENTA.

PIGMENTUM ACIDI SULPHUROSI.

(Throat).

Sulphurous Acid	2 fluid drachms.
Glycerin of Tannic Acid ...	2 fluid drachms.
Glycerin	4 fluid drachms.

PIGMENTUM ARGENTI NITRATIS ÆTHEREUM.

(Throat).

Silver Nitrate	... 20 grains.
Distilled Water ...	... 1 fluid drachm.
Spirit of Nitrous Ether	... to 1 fluid ounce.

Pigmentum Chloral Compositum.

Chloral Hydrate	of each equal parts.
Menthol	
Camphor	

PIGMENTUM FERRI PERCHLORIDI.

(Throat).

Solution of Ferric Chloride...	of each equal parts.
Glycerin	

Pigmentum Hydrargyri Nitratis.

Nitrate of Mercury Ointment	... 40 grains.
Olive Oil	½ fluid ounce.
Liquid Paraffin	to 1 fluid ounce.

Pigmentum Iodi.

Iodine...	... 30 grains.
Methylated Spirit	to 1 fluid ounce.

PIGMENTUM IODI CARBOLIZATUM.

(Obstetric).

Iodine	1 part.
Liquefied Carbolic Acid	4 parts.

PIGMENTUM IODI COMPOSITUM.

(Throat).

Iodine	5 grains.
Potassium Iodide	5 grains.
Oil of Peppermint	5 minims.
Glycerin	to 1 fluid ounce.

PILULÆ (*See also Tabellæ*).

Pilula Ferri et Arsenici.

Arsenious Anhydride	$\frac{1}{36}$ grain.
Iron Pill	...to 5 grains.

Pilula Ferri et Arsenici cum Quinina.

Arsenious Anhydride	$\frac{1}{36}$ grain.
Quinine Sulphate	1 grain.
Iron Pill	...to 5 grains.

Pilula Ferri et Arsenici cum Strychnina.

Arsenious Anhydride	$\frac{1}{36}$ grain.
Strychnine	$\frac{1}{60}$ grain.
Iron Pill	...to 5 grains.

Pilula Ferri et Quininæ cum Strychnina.

Quinine Sulphate	1 grain.
Strychnine	$\frac{1}{60}$ grain.
Iron Pill	...to 5 grains.

Pilula Ferri et Strychninæ.

Strychnine	$\frac{1}{60}$ grain.
Iron Pill	...to 5 grains.

POTUS.

Potus Imperialis.

Acid Potassium Tartrate	$\frac{1}{4}$ ounce.
Lemon Juice	$\frac{1}{2}$ fluid ounce.
Syrup	$\frac{1}{2}$ fluid ounce.
Water	to 1 pint.

Note :—The Syrup should be omitted for Diabetic patients.

PULVERES.

Pulvis Acidi Salicylici Compositus.

Dusting Powder.

Salicylic Acid in powder	3 parts.
French Chalk in powder	87 parts.
Lycopodium	10 parts.

Pulvis Bismuthi Compositus.

Syn : **Anti-Dyspeptic Powder.**

Bismuth Oxycarbonate ...	of each equal parts.
Heavy Magnesium Carbonate	
Sodium Bicarbonate... ...	

Dose :—1 drachm.

Pulvis Bismuthi cum Morphina.

Syn : **Ferrier's Snuff.**

Bismuth Oxynitrate	... 6 drachms.
Morphine Hydrochloride ...	... 2 grains.
Gum Acacia in powder ...	... 2 drachms.

Dose for Insufflation :—120 grains during 24 hours.

Pulvis Guaiaci Compositus.

Guaiacum Resin in powder... ...	1 part.
Heavy Magnesium Carbonate ...	2 parts.
Sublimed Sulphur	1 part.

Dose :—15 to 30 grains.

Pulvis Iodoformi et Acidi Borici.

Dusting Powder.

Iodoform in powder	1 part.
Boric Acid in powder	3 parts.

Pulvis Iodoformi et Morphinæ.

Iodoform in powder	30 grains.
Morphine Hydrochloride	3 grains.
Boric Acid in powder	87 grains.

Dose for Laryngeal Insufflation :—1 to 2 grains.

Pulvis Lobeliæ Compositus.

Lobelia in powder	of each equal parts.
Anise Fruit in powder ...	
Potassium Nitrate in powder	
Stramonium Leaves in powder	

Dose for Inhalation :—The fumes of 60 grains of the burnt powder.

Pulvis Mentholis Compositus.

Menthol in powder	2 drachms.
Ammonium Chloride in powder ...	2 drachms.
Boric Acid in powder	4 drachms.

Dose for insufflation :—60 grains during 24 hours.

Pulvis Sodii Sulphatis Compositus.

Syn: **Carlsbad Salt.**

Exsiccated Sodium Sulphate ...	22 parts.
Potassium Sulphate	1 part.
Sodium Bicarbonate	18 parts.
Sodium Chloride	9 parts.

Dose:—1 drachm in Warm Water.

Pulvis Zinci Oxidi et Acidi Borici.

Dusting Powder.

Zinc Oxide	of each equal parts.
Boric Acid in powder ...	

Pulvis Zinci Oxidi et Calomelanos.

Dusting Powder.

Zinc Oxide	3 parts.
Calomel	1 part.

SUPPOSITORIA.

Suppositorium Cocainæ.

Cocaine	¼ or ½ grain.
Oil of Theobroma	to 15 grains.

Suppositorium Hamamelidis.

Liquid Extract of Hamamelis ...	5 minims.
Oil of Theobroma	to 15 grains.

SYRUPI.

Syrupus Ferri Phosphatis Compositus.

Syn.: **Chemical Food.**

Contains in each fluid drachm about ½ grain of Iron Phosphate, 1 grain of Calcium Phosphate, with small quantities of Potassium and Sodium Phosphates.

Syrupus Hypophosphitum Compositus.

Contains in each fluid drachm 1 grain of Iron Hypophosphite, ½ grain of Manganese Hypophosphite, 1 grain of Calcium Hypophosphite and ½ grain of Potassium Hypophosphite.

TABELLÆ.

Tabella Aloini Composita.

Aloin	of each ½ grain.
Extract of Nux Vomica	
Ferrous Sulphate	
Hard Soap in powder	
Myrrh in powder	

Tabella Bismuthi et Calomelanos pro Infantibus.

Bismuth Oxycarbonate	5 grains.
Calomel	⅙ grain.
Compound Powder of Ipecacuanha	⅙ grain.

Tabella Bismuthi Composita pro Infantibus.

Bismuth Oxynitrate	5 grains.
Aromatic Powder of Chalk ...	2 grains.
Heavy Magnesium Carbonate ...	2 grains.

Tabella Calomelanos Composita.

Calomel	$\frac{1}{4}$ grain.
Green Extract of Belladonna ...	$\frac{1}{8}$ grain.
Quinine Sulphate	1 grain.

Tabella Calomelanos cum Opio.

Calomel	1 grain.
Opium in powder	$\frac{1}{4}$ grain.

Tabella Cascaræ Composita.

Extract of Cascara Sagrada ...	1 grain.
Aloin	$\frac{1}{2}$ grain.
Extract of Hyoscyamus	1 grain.
Extract of Nux Vomica	$\frac{1}{4}$ grain.

Tabella Cathartica Composita.

Calomel	$\frac{1}{2}$ grain.
Jalap in powder	3 grains.
Socotrine Aloes	1 grain.

Tabella Colocynthidis et Calomelanos.

Compound Pill of Colocynth ...	4 grains.
Calomel	1 grain.

Tabella Colocynthidis et Hydrargyri cum Creta.

Compound Pill of Colocynth ...	3 grains.
Mercury with Chalk...	2 grains.

Tabella Digitalis et Hydrargyri cum Creta.

Digitalis Leaf in powder	of each 1 grain.
Mercury with Chalk	
Squill in powder	

Tabella Ferri Aperiens.

Ferrous Sulphate	1 grain.
Extract of Nux Vomica	$\frac{1}{4}$ grain.
Socotrine Aloes	1 grain.

Tabella Ferri Phosphatis cum Quinina et Strychnina.

Iron Phosphate	$\frac{1}{2}$ grain.
Quinine Sulphate	$\frac{2}{5}$ grain.
Strychnine	$\frac{1}{64}$ grain.

Tabella Ferri Redacti et Hydrargyri cum Creta.

Reduced Iron	2 grains.
Mercury with Chalk	1 grain.

Tabella Ferri Sulphatis cum Strychnina.

Ferrous Sulphate	3 grains.
Strychnine	$\frac{1}{32}$ grain.

Tabella Hydrargyri Composita.

Mercury with Chalk	1$\frac{1}{2}$ grains.
Extract of Colchicum	$\frac{1}{4}$ grain.
Ipecacuanha Root in powder ...	$\frac{1}{4}$ grain.
Pill of Colocynth and Hyoscyamus..	3 grains.

Tabella Hydrargyri cum Creta et Rheo.

Mercury with Chalk	2½ grains.
Compound Rhubarb Pill	2½ grains.

Tabella Hydrargyri cum Creta et Rheo pro Infantibus.

Mercury with Chalk	½ grain.
Rhubarb in powder	2½ grains.
Ginger in powder	½ grain.
Sodium Bicarbonate	1 grain.

Tabella Hydrargyri cum Creta et Soda pro Infantibus.

Mercury with Chalk	½ grain.
Sodium Bicarbonate	2 grains.

Tabella Hydrargyri et Ipecacuanhæ Composita.

Mercury with Chalk	1 grain.
Compound Powder of Ipecacuanha ..	1 grain.

Tabella Nucis Vomicæ Composita.

Extract of Nux Vomica	⅙ grain.
Alcoholic Extract of Belladonna ...	¼ grain.
Aloin	½ grain.

Tabella Phenacetini et Caffeinæ Citratis.

Phenacetine	5 grains.
Caffeine Citrate	3 grains.

Tabella Rhei et Sodæ pro Infantibus.

Rhubarb in powder	3 grains.
Sodium Bicarbonate	1 grain.

Tabella Santonini et Calomelanos pro Infantibus.

Santonin	1 grain.
Calomel	$\frac{1}{2}$ grain.
Refined Sugar in powder	$1\frac{1}{2}$ grains.

Tabella Santonini Composita pro Infantibus.

Santonin	1 grain.
Calomel	$\frac{1}{2}$ grain.
Compound Powder of Scammony ...	$2\frac{1}{2}$ grains.

Tabella Scammonii Composita pro Infantibus.

Scammony Resin in powder ...	1 grain.
Acid Potassium Tartrate	1 grain.
Antimonial Powder	$\frac{1}{3}$ grain.
Calomel	1 grain.
Ginger in powder	$\frac{1}{3}$ grain.
Jalap in powder	$\frac{1}{3}$ grain.

Tabella Zinci Oxidi et Belladonnæ.

Zinc Oxide	2 grains.
Green Extract of Belladonna ...	$\frac{1}{4}$ grain.

The following Drugs are also kept in Tablet Form in the various strengths mentioned:—

Acidum Acetyl-Salicylicum (Syn: Aspirin)	5, 10 grains.
Acidum Arseniosum	$\frac{1}{30}$ grain.
Acidum Carbolicum	$\frac{1}{2}$, 1 grain.
Acidum Diethylbarbituricum (Syn: Veronal)	$3\frac{3}{4}$ grains.

Drugs in Tablet Form (*continued*) :—

Ammonii Sulpho-ichthyolas (Syn: Ichthyol)	2 grains.
Caffeine Citras	2½, 5 grains.
Calomel	⅙, ¼, ½, 1, 2, 3, 5 grains.
Cerii Oxalas	3 grains.
Codeina	¼, ⅓, ½, 1, 2, 3 grains.
Creosotum	1, 2 minims.
Extractum Belladonnæ Alcoholicum	¼, ⅓, ½ grain.
Extractum Cannabis Indicæ ...	¼, ½ grain.
Extractum Cascaræ Sagradæ ...	2, 4 grains.
Extractum Ergotæ	2 grains.
Ferri Sulphas...	2, 3 grains.
Ferrum Redactum	3, 5 grains.
Glusidum	½ grain.
Hydrargyrum cum Creta	¼, ½, 1, 2, 3 grains.
Methyl Sulphonal (Syn: Trional)	5, 10 grains.
Morphinæ Hydrochloridum...	¼, ⅓, ½ grain.
Opium...	¼, ⅓, ½, 1 grain.
Pepsinum	2½ grains.
Phenacetinum	5, 10 grains.
Phenazonum	5, 10 grains.
Phenolphthaleïn	¾, 1½, 7 grains.
Podophylli Resina	⅙, ¼, ½ grain.
Pulvis Ipecacuanhæ Compositus	5, 10 grains.
Quininæ Sulphas	1, 2, 3, 5 grains.
Salicinum	10 grains.
Salol	3, 5, 10 grains.
Sulphonal	5, 10 grains.
Suprarenal Gland, fresh substance...	5 grains.
Theobrominum Sodio - Salicylicum (Syn: Diuretin)	5 grains.
Thyroideum Siccum... ...	1, 3, 5 grains.
Zinci Oxidum	2½, 5 grains.
Zinci Valerianas	...2, 3 grains.

TROCHISCUS.

Trochiscus Glycyrrhizæ.

Extract of Liquorice	...	3 grains.
Oil of Anise	...	½ minim.
Refined Sugar ...	...	to 15 grains.

UNGUENTA.

UNGUENTUM ACIDI BORICI.

(Ophthalmic).

Boric Acid in powder	...	1 drachm.
Soft Paraffin	...	... to 1 ounce.

Unguentum Acidi Carbolici Compositum.

Carbolic Acid	...	1 part.
Mercuric Nitrate Ointment	...	2 parts.
Sulphur Ointment	...	2 parts.

UNGUENTUM ATROPINÆ.

(Ophthalmic).

Atropine Sulphate ...	...	4 grains.
Soft Paraffin	...	... to 1 ounce.

UNGUENTUM ATROPINÆ FORTE.

(Ophthalmic).

Atropine Sulphate ...	...	8 grains.
Soft Paraffin	...	... to 1 ounce.

Unguentum Chrysarobini Compositum.

Chrysarobin	5 grains.
Ammoniated Mercury	10 grains.
Solution of Coal Tar	10 minims.
Soft Paraffin	to 1 ounce.

UNGUENTUM COCAINÆ.

(Ophthalmic).

Cocaine Hydrochloride	8 grains.
Soft Paraffin	to 1 ounce.

Unguentum Cupri Oleatis.

Copper Oleate	1 part.
Soft Paraffin	7 parts.

Unguentum Diachyli.

Lead Plaster ... ... }	of each equal
Soft Paraffin ... ... }	parts.

UNGUENTUM ESERINÆ.

(Ophthalmic).

Eserine Sulphate	1 grain.
Soft Paraffin	to 1 ounce.

UNGUENTUM HYDRARGYRI AMMONIATI DILUTUM.

(Ophthalmic).

Ammoniated Mercury	8 grains.
Soft Paraffin	to 1 ounce.

Unguentum Hydrargyri Ammoniati Dilutum.

Ammoniated Mercury Ointment ...	1 part.
Soft Paraffin	3 parts.

UNGUENTUM HYDRARGYRI OXIDI FLAVI.

(Ophthalmic).

Yellow Mercuric Oxide	4 grains.
Soft Paraffin	... to 1 ounce.

UNGUENTUM HYDRARGYRI OXIDI FLAVI FORTE.

(Ophthalmic).

Yellow Mercuric Oxide	8 grains.
Soft Paraffin	... to 1 ounce.

UNGUENTUM HYDRARGYRI OXIDI FLAVI CUM ATROPINA.

(Ophthalmic).

Yellow Mercuric Oxide	4 grains.
Atropine Sulphate	2 grains.
Soft Paraffin	... to 1 ounce.

UNGUENTUM HYDRARGYRI OXIDI FLAVI CUM ATROPINA FORTE.

(Ophthalmic).

Yellow Mercuric Oxide	8 grains.
Atropine Sulphate	4 grains.
Soft Paraffin	... to 1 ounce.

UNGUENTUM HYDRARGYRI OXIDI FLAVI CUM ESERINA.

(Ophthalmic).

Yellow Mercuric Oxide	4 grains.
Eserine Sulphate	1 grain.
Soft Paraffin	... to 1 ounce.

Unguentum Hydrargyri Oxidi Rubri Dilutum.

Red Mercuric Oxide Ointment ...	1 part.
Soft Paraffin	7 parts.

Unguentum Hydrargyri Perchloridi Compositum.

Mercuric Chloride	2 grains.
Carbolic Acid	20 grains.
Glycerin	10 minims.
Olive Oil	40 minims.
Zinc Ointment	to 1 ounce.

Unguentum Ichthyolis.

Ammonium Sulpho-ichthyolate ...	40 grains.
Salicylic Acid in powder	8 grains.
Soft Paraffin	to 1 ounce.

Unguentum Metallorum.

Calomel	10 grains.
Lead Acetate in powder	10 grains.
Zinc Oxide	20 grains.
Diluted Mercuric Nitrate Ointment	to 1 ounce.

Unguentum Naphtholis Compositum.

Naphthol in powder...	1½ parts.
Prepared Chalk in powder	1 part.
Soft Soap (B.P.)	5 parts.
Lard	10 parts.

Unguentum Olei Cadini.

Oil of Cade	1 fluid drachm.
Soft Paraffin	... to 1 ounce.

Unguentum Picis Carbonis cum Hydrargyro Ammoniato.

Solution of Coal Tar ...	... 30 minims.
Ammoniated Mercury ...	... 10 grains.
Soft Paraffin	... to 1 ounce.

Unguentum Pyrogallolis Compositum.

Pyrogallol	... 20 grains.
Ammonium Sulpho-ichthyolate	... 20 grains.
Salicylic Acid in powder	... 8 grains.
Soft Paraffin	... to 1 ounce.

Unguentum Resorcini.

Resorcin	... 1 drachm.
Glycerin	1 fluid drachm.
Hydrous Wool Fat ...	... 2 drachms.
Soft Paraffin	... to 1 ounce.

Unguentum Sulphuris et Zinci Oxidi.

Sulphur Ointment ...	1 part.
Zinc Ointment ...	7 parts.

Unguentum Thymolis.

Thymol	... 20 grains.
Soft Paraffin	... to 1 ounce.

Unguentum Zinci Carbonatis et Adipis Lanæ Hydrosi.

Zinc Carbonate ...	... 1 drachm.
Salicylic Acid in powder	... 8 grains.
Hydrous Wool Fat ...	... 2 drachms.
Soft Paraffin	... to 1 ounce.

Unguentum Zinci Oxidi et Adipis Lanæ Hydrosi.

Zinc Oxide	1 drachm.
Salicylic Acid in powder	10 grains.
Hydrous Wool Fat	2 drachms.
Soft Paraffin	to 1 ounce.

Unguentum Zinci Oxidi cum Resorcino.

Resorcin	15 grains.
Zinc Ointment	to 1 ounce.

Unguentum Zinci Peroxidi.

Zinc Peroxide	20 grains.
Soft Paraffin	to 1 ounce.

Unguentum Zinci Peroxidi Forte.

Zinc Peroxide	40 grains.
Soft Paraffin	to 1 ounce.

VAPORES.

Vapor Acidi Carbolici.

Liquefied Carbolic Acid ...	30 minims.
Boiling Water	1 pint.

Vapor Benzoini.

Compound Tincture of Benzoin	1 fluid drachm.
Boiling Water	1 pint.

Vapor Creosoti.

Creosote	... 10 minims.
Boiling Water	... 1 pint.

Vapor Cubebæ.

Oil of Cubebs	... 5 minims.
Boiling Water	... 1 pint.

Vapor Eucalypti.

Oil of Eucalyptus	... 5 minims.
Boiling Water	... 1 pint.

Vapor Iodi.

Strong Solution of Iodine ...	... 10 minims.
Boiling Water	... 1 pint.

Vapor Mentholis.

Solution of Menthol (1 in 3)	... 5 minims.
Boiling Water	... 1 pint.

Vapor Pini Sylvestris.

Oil of Scotch Pine	... 5 minims.
Boiling Water	... 1 pint.

Vapor Terebeni.

Terebene	... 5 minims.
Boiling Water	... 1 pint.

Vapor Terebinthinæ.

Turpentine	1 fluid drachm.
Boiling Water	... 1 pint.

Beef Juice.

Scrape ¼ lb. Lean Raw Beef into a small basin, cover with Cold Water and set aside for an hour in a cool place, protected from dust. Then forcibly compress with the hand and strain through muslin. The product should measure 4 ounces and contain 5% Coagulable Protein.

Beef Tea.

Take 1 lb. of Fresh Lean Beef, remove all Gristle and Fat and cut it in small pieces, or scrape down thoroughly into shreds with the back of a knife. Place in a stone jar, having a tightly fitting lid, and add half a teaspoonful of Salt and one pint of Cold Water. Allow the Meat to soak for ten minutes, stirring occasionally. Then place the jar in a saucepan half full of Hot Water and maintain at a temperature of about 160°F (71°C), *i.e.*, below the coagulating point of meat. Stir the mixture from time to time with two forks, separating the cakes of Meat and squeezing them against the sides of the jar. At the end of an hour raise the temperature to boiling point for a moment and pour off the Tea from the residue immediately, holding back and squeezing the pieces of Meat with a fork to extract the juice. Set aside till cold and remove the accumulated Fat with a heated spoon. The product should measure about a pint and contain at least 1½% of Protein. The flavour may be improved by adding towards the end of the process a little Parsley, or Thyme and Pay Leaf tied together, or any other palatable herb or vegetable.

Whole Beef Tea may be made by passing some of the fibre of the Meat several times through a

mincing machine and adding it to the Tea. The nutritive value of the latter is greatly enhanced by this addition.

Egg White Solution.

Take the White of a Raw Egg and cut it in small portions with a clean pair of scissors. Place in a suitable vessel and add a pinch of Salt and half a pint of Cold Water. Then shake and strain through muslin. A small quantity of White Sugar or Liebig's Extract, dissolved in a little Warm Water, may be used instead of Salt as a flavouring. The product will contain $1\frac{1}{4}$ % Coagulable Protein.

Normal Saline.

To 2 pints of Cold, Sterilized, Concentrated Liquor, containing 240 grains of Common Salt, add 1 pint of Boiling Water, and a solution of required strength and temperature will be obtained ready for use.

Peptonised Milk.

Dilute a pint of Milk with a quarter of a pint of Water and heat to a temperature of about 120°F (49°C). Add 1 Zymine Powder, or 20 grains of Sodium Bicarbonate and 2 teaspoonfuls of Liquor Pancreaticus. Then pour into a jar provided with a cover, and set aside in a warm place for one to one and a half hours. Finally bring to boil and cool rapidly.

DIET WITH MINIMUM QUANTITY OF CARBOHYDRATES.

—o—

Breakfast.

Bacon and Scrambled Eggs, Casein Bread (such as Protene or that prepared by Callard & Co. is recommended) toasted, with plenty of Butter, Tea or Coffee, with Sugar Free Milk and Saccharin.

Between Breakfast and Dinner, ½ pint of Sugar Free Milk.

Dinner.

Fish, Meat (6 ounces), Green Vegetables or Turnips with Butter, Custard of ½ to 1 pint Sugar Free Milk and Eggs, Whisky (2 fluid ounces).

Tea.

One or two Eggs or Cheese, Casein Bread, and Butter, Tea or Coffee, with Sugar Free Milk and Saccharin.

Supper.

Fish (2 ounces) with Melted Butter, Casein Bread, and 1 pint of Sugar Free Milk.

A strict diet for Diabetics should only be adopted in those cases in which marked Acetonuria has been proved to be absent. This is shown by the addition of Ferric Chloride to the Urine failing to produce a port wine colour.

If a patient suffering from marked Acetonuria be placed suddenly upon a strict diet, Coma not infrequently results.

INFANT FEEDING.

—o—

The following table gives the quantities suitable for a healthy child :—

Age of Child.	Milk.	Water or Barley Water.	Total amount to be given at each meal.
During—	Tablespoons.	Tablespoons.	Tablespoons.
1st fortnight	1	2	3
2nd ,,	2	3	5
2nd month	2	3	5
3rd ,,	4	4	8
4th ,,	5	4	9
5th ,,	6	4	10
6th ,,	8	4	12
7th ,,	9	4	13
8th ,,	10	4	14
9th ,,	12	4	16

If the child does not thrive on Milk alone, a small quantity of Fresh Cream may be added to each feed.

Children may be fed with a spoon or from a bottle; the latter must be without any tubes, and should have the teat fitting directly upon it.

The Boat Bottle marked in Tablespoons is very suitable.

Bottles and Teats must be washed once a day at least with boiling water, to which a little Washing Soda has been added, and again rinsed with cold water.

They should always be rinsed after each feed and kept under cold water.

After eight months of age, and if teeth are present, the child may have Bread and Milk, Bread and Butter, Milk Pudding, a little Red Gravy with Bread Crumbs, or a little Mashed Potato, or the Yolk of a slightly boiled Egg, once a day, at the mid-day meal.

It is advisable to vary the diet occasionally.

After nine months, Eggs, Mutton Broth, or Bread fried in Bacon Fat, may be gradually added to the diet.

Fish or Meat should not be given before the child is eighteen months old; when given, it should be finely minced.

Children, especially those under three years of age, should not be fed upon "just what the parents have."

Tea or Coffee should not be given to children under five years of age.

Stimulants should never be given to children, except by the direction of a medical man.

DIRECTIONS FOR THE CARE OF RICKETY CHILDREN.

—o—

Rickets is a disease which shows its effects principally in the Bones of the Limbs, but which, nevertheless, affects all other parts of the body.

To improve the condition the greatest care is necessary in giving the child suitable food.

The Diet must consist of Cow's Milk and things made with it, Bread and Milk, Milk Puddings, etc., and when the child is more than a year old, Gravy, to which Scalded Bread may be added, Cooked Vegetables, Eggs slightly boiled or beaten up in Milk, and a little Bread and Butter.

The child must be fed at regular times and given nothing between meals.

It must *not* be given Sweets, Biscuits, Buns, Cakes, Pastry, or patent foods of any kind.

The child should not be clothed too warmly by night or day. It should be taken out of doors as much as po sible, when the weather is fine, and the room in which it sleeps must be kept fresh and cool.

If Splints are ordered for the child's legs, be careful to observe how they are put on. They must be removed in the evening when the child is put to bed, and put on again in the morning, the ends of the splints projecting beyond the feet, and the bandages applied right up to the top of the thighs, so that the child cannot walk, stand, or double its legs up when sitting.

THERMOMETRIC TABLE.

Comparison of Fahrenheit and Centigrade Scales.





WEIGHTS AND MEASURES OF THE IMPERIAL SYSTEM.

—o—

Measures of Mass.

1 Grain	gr.		
1 Ounce (Avoir.)	oz.		= 437·5 grains.
1 Pound	lb.	= 16 ounces	= 7000 grains.

Measures of Capacity.

1 Minim	min.		
1 Fluid Drachm	fl. drm.	=	60 minims.
1 Fluid Ounce	fl. oz.	=	8 fluid drachms.
1 Pint	O.	=	20 fluid ounces.
1 Gallon	C.	=	8 pints.

Measures of Length.

1 Inch	in.		
1 Foot	ft.	=	12 inches.
1 Yard	yd.	=	36 inches.

Relation of Volume to Mass.

1 Minim is the volume at 62° F. of ·911 grain of water.

1 Fluid Drachm is the volume at 62° F. of 54·687 grains of water.

1 Fluid Ounce is the volume of 1 ounce or 437·5 grains of water.

1 Pint is the volume of 1·25 pounds or 8750 grains of water.

1 Gallon is the volume of 10 pounds or 70000 grains of water.

110 Minims nearly = the volume at 62° F. of 100 grains of water.

WEIGHTS AND MEASURES OF THE METRIC SYSTEM.

—o—

Measures of Mass.

1 Milligramme = the thousandth part of one grm. or 0·001 grm.

1 Centigramme = the hundredth part of one grm. or 0·01 grm.

1 Decigramme = the tenth part of 1 grm. or 0·1 grm.

1 Gramme = weight of one millilitre of distilled water at 4° C. (39·2° F.) 1·0 grm.

1 Dekagramme = ten grammes or 10·0 grm.

1 Hectogramme = one hundred grammes or 100·0 grm.

1 Kilogramme = one thousand grammes or 1000·0 grm.

Measures of Capacity.

1 Millilitre or 1 Cubic Centimetre = the volume at 4° C. of 1 grm. of water.

1 Centilitre = the volume at 4° C. of 10 grms. of water.

1 Decilitre = the volume at 4° C. of 100 grms. of water.

1 Litre = the volume at 4° C. of 1000 grms. (1 kilog.)

Measures of Length.

1 Millimetre = one thousandth part of one metre or 0·001 metre.

1 Centimetre = one hundredth part of one metre or 0·01 metre.

1 Decimetre = one tenth part of one metre or 0·1 metre.

1 Metre = 1·0 metre

Relation of the Imperial Standards to the Metric Standards.

Standards of Mass.

1 Grain = 0·064 gramme nearly.
1 Ounce = 28·349 grammes nearly.
1 Pound = 453·592 grammes.

Standards of Capacity.

1 Minim = 0·059 cubic centimetre nearly.
1 Fluid Drachm = 3·552 cubic centimetres nearly.
1 Fluid Ounce = 28·417 ,, ,, ,,
1 Pint = 568·336 cubic centimetres nearly.
1 Gallon = 4·545 litres.

Standards of Length.

1 Inch = 2·51 centimetres.
1 Foot = 30·48 ,,
1 Yard = 91·44 ,,

Relation of the Metric Standards to the Imperial Standards.

Standards of Mass.

1 Milligramme = 0·015 grain nearly.
1 Centigramme = 0·154 ,, ,,
1 Decigramme = 1·543 grains ,,
1 Gramme = 15·432 grains.
1 Kilogramme = 35 ounces 120 gr. or 15432 grains.

Standards of Capacity.

1 Cubic Centimetre = 17 minims nearly.
1 Litre = 35 fl. oz. 1 fl. dr. 34 min. nearly.

Standards of Length.

1 Millimetre = 0·039 inch.
1 Centimetre = 0·393 ,,
1 Decimetre = 3·937 inches.
1 Metre = 39·37 inches nearly.

Relation of the Imperial Standards to the Metric Standards.

STANDARDS OF MASS.

1 Grain = 0·004 gramme nearly.
1 Ounce = 28·349 grammes nearly.
1 Pound = 453·592 grammes.

STANDARDS OF CAPACITY.

1 Minim = 0·059 cubic centimetre nearly.
1 Fluid Drachm = 3·552 cubic centimetres nearly.
1 Fluid Ounce = 28·412 " "
1 Pint = 568·239 cubic centimetres nearly.
1 Gallon = 4·543 litres.

STANDARDS OF LENGTH.

1 Inch = 2·54 centimetres.
1 Foot = 30·48 "
1 Yard = 91·44 "

Relation of the Metric Standards to the Imperial Standards.

STANDARDS OF MASS.

1 Milligramme = 0·015 grain nearly.
1 Centigramme = 0·154 "
1 Decigramme = 1·543 grains "
1 Gramme = 15·432 grains
1 Kilogramme = 35 ounces 120 gr. or 15432 grains.

STANDARDS OF CAPACITY.

1 Cubic Centimetre = 17 minims nearly.
1 Litre = 35 fl. oz. 1 fldr. 34 minims nearly.

STANDARDS OF LENGTH.

1 Millimetre = 0·039 inch.
1 Centimetre = 0·393 "
1 Decimetre = 3·937 inches.
1 Metre = 39·37 inches nearly.

POSOLOGICAL TABLE.

Name.	Dose.

[The various forms in which certain drugs are now prepared for internal use will be found on pages indicated by numerals in brackets].

Acetanilidum	1 to 3 gr.
Acetum Ipecacuanhæ...	10 to 30 min.
— Scillæ (62)	10 to 30 min.
Acidum Aceticum Dilutum (58) ...	½ to 2 fl. dr.
— Acetyl-Salicylicum (75)	10 to 15 gr.
— Arseniosum (67, 75)	$\frac{1}{60}$ to $\frac{1}{15}$ gr.
— Benzoicum	5 to 15 gr.
— Boricum	5 to 15 gr.
— Carbolicum (75)	1 to 3 gr.
— — Liquefactum	1 to 3 min.
— Citricum (61)	5 to 20 gr.
— Diethylbarbituricum (75)... ...	5 to 15 gr.
— Gallicum	5 to 15 gr.
— Hydrobromicum Dilutum (61) ...	15 to 60 min.
— Hydrochloricum Dilutum (57) ...	5 to 20 min.
— Hydrocyanicum Dilutum	2 to 6 min.
— Nitricum Dilutum	5 to 20 min.
— Nitro-hydrochloricum Dilutum ...	5 to 20 min.
— Phosphoricum Dilutum	5 to 20 min.
— Salicylicum...	5 to 20 gr.
— Sulphuricum Aromaticum (53) ...	5 to 20 min.
— — Dilutum (58)	5 to 20 min.
— Sulphurosum	½ to 1 fl. dr.
— Tannicum	2 to 5 gr.
— Tartaricum	5 to 20 gr.
Æther	10 to 30 min. for repeated doses; for a single dose, 40 to 60 min.

Name.	Dose.
Æther Aceticus	20 to 40 min. for repeated doses; for a single dose, 60 to 90 min.
Aloe Barbadensis	2 to 5 gr.
— Socotrina (72, 73)	2 to 5 gr.
Aloinum (71, 72, 74)	½ to 2 gr.
Alumen	5 to 10 gr.
Ammoniacum	5 to 15 gr.
Ammonii Benzoas	5 to 15 gr.
— Bromidum	5 to 30 gr.
— Carbonas (54, 61, 63)	3 to 10 gr.
— Chloridum	5 to 20 gr.
— Phosphas	5 to 20 gr.
— Sulphoichthyolas (76) ...	10 to 30 gr.
Amyl Nitris	For inhalation—The vapour of 2 to 5 min.
Antimonii Oxidum	1 to 2 gr.
Antimonium Sulphuratum	1 to 2 gr.
— Tartaratum	Diaphoretic, $\frac{1}{24}$ to $\frac{1}{8}$ gr.; emetic, 1 to 2 gr.
Apomorphinæ Hydrochloridum ...	$\frac{1}{20}$ to $\frac{1}{10}$ gr. by subcutaneous injection; by the mouth, $\frac{1}{10}$ to $\frac{1}{4}$ gr.
Aqua Laurocerasi (47)	½ to 2 fl. dr.
Argenti Nitras	$\frac{1}{4}$ to ½ gr.
— Oxidum	½ to 2 gr.

Name.	Dose.
Arsenii Iodidum	$\frac{1}{20}$ to $\frac{1}{5}$ gr.
Asafetida	5 to 15 gr.
Atropina	$\frac{1}{200}$ to $\frac{1}{100}$ gr.
Atropinæ Sulphas	$\frac{1}{200}$ to $\frac{1}{100}$ gr.
Balsamum Peruvianum	5 to 15 min.
— Tolutanum	5 to 15 gr.
Bismuthi Oxidum	5 to 20 gr.
— Oxycarbonas (54, 68, 71)	5 to 20 gr.
— Oxynitras (72)	5 to 20 gr.
— Salicylas	5 to 20 gr.
Borax	5 to 20 gr.
Butyl-Chloral Hydras	5 to 20 gr.
Caffeina	1 to 5 gr.
Caffeinæ Citras (74, 76)	2 to 10 gr.
— — Effervescens	60 to 120 gr.
— Sodio-Salicylas	1 to 4 gr., hypodermically.
Calcii Carbonas Præcipitatus	10 to 60 gr.
— Chloridum	5 to 15 gr.
— Hypophosphis (71)	3 to 10 gr.
— Phosphas (71)	5 to 15 gr.
Calomel (71, 75, 76)	$\frac{1}{2}$ to 5 gr.
Calx Sulphurata	$\frac{1}{4}$ to 1 gr.
Cambogia	$\frac{1}{2}$ to 2 gr.
Camphora	2 to 5 gr.
Carbo Ligni	60 to 120 gr.
Catechu	5 to 15 gr.
Cerii Oxalas (76)	2 to 10 gr.
Chloral Hydras (44)	5 to 20 gr.
Chloroformum (35)	1 to 5 min.
Cocainæ Hydrochloridum	$\frac{1}{5}$ to $\frac{1}{2}$ gr.

Name.	Dose.
Codeina (76)	¼ to 2 gr.
Codeinæ Phosphas	¼ to 2 gr.
Colchici Cormus	2 to 5 gr.
Confectio Piperis (34)	60 to 120 gr.
— Sennæ (34)	60 to 120 gr.
— Sulphuris (34)	60 to 120 gr.
Copaiba	½ to 1 fl. dr.
Creosotum (76)	1 to 5 min.
Creta Præparata	10 to 60 gr.
Cubebæ Fructus	30 to 60 gr.
Cupri Sulphas	¼ to 2 gr. as an astringent; as an emetic, 5 to 10 gr.
Cusso	¼ to ½ oz.
Decoctum Aloes Compositum (57, 58)	½ to 2 fl. oz.
— Cinchonæ (56, 57, 61)	1 to 2 fl. oz.
— Granati Corticis	½ to 2 fl. oz.
— Hæmatoxyli	½ to 2 fl. oz.
Diacetyl-Morphinæ Hydrochloridum (47)	$\frac{1}{12}$ to ⅙ gr.
Digitalis Folia (73)	½ to 2 gr.
Diuretin (76)	5 to 15 gr.
Elaterinum	$\frac{1}{40}$ to $\frac{1}{10}$ gr.
Elaterium	$\frac{1}{10}$ to ½ gr.
Ergota	20 to 60 gr.
Ergotinæ Citras	$\frac{1}{150}$ to $\frac{1}{30}$ gr.
Eucalypti Gummi	2 to 5 gr.
Extractum Aloes Barbadensis	2 to 4 gr.
— Anthemidis	2 to 8 gr.
— Belladonnæ Alcoholicum (74, 76)	¼ to 1 gr.
— — Viride (72, 75)	¼ to 1 gr.
— Cannabis Indicæ (76)	¼ to 1 gr.

Name.	Dose.
Extractum Cascaræ Sagradæ (72, 76) ...	2 to 8 gr.
— — — Liquidum (54, 56, 57) ...	½ to 1 fl. dr.
— Cimicifugæ Liquidum	5 to 30 min.
— Cinchonæ Liquidum	5 to 15 min.
— Cocæ Liquidum	½ to 1 fl. dr.
— Colchici (73)	¼ to 1 gr.
— Colocynthidis Compositum ...	2 to 8 gr.
— Ergotæ (76)	2 to 8 gr.
— — Liquidum (57)	10 to 30 min.
— Euonymi Siccum	1 to 2 gr.
— Filicis Liquidum (44)	45 to 90 min.
— Gentianæ	2 to 8 gr.
— Glycyrrhizæ (77)	2 to 8 gr.
— — Liquidum (44, 56, 57)	½ to 1 fl. dr.
— Hamamelidis Liquidum	5 to 15 min.
— Hydrastis Liquidum	5 to 15 min.
— Hyoscyami Viride (72)	2 to 8 gr.
— Ipecacuanhæ Liquidum	Expectorant, ½ to 2 min.; emetic, 15 to 20 min.
— Jaborandi Liquidum	5 to 15 min.
— Jalapæ	2 to 8 gr.
— Krameriæ	5 to 15 gr.
— Nucis Vomicæ (71, 72, 73, 74) ...	¼ to 1 gr.
— — — Liquidum	1 to 3 min.
— Opii	¼ to 1 gr.
— — Liquidum	5 to 30 min.
— Pareiræ Liquidum	½ to 2 fl. dr.
— Physostigmatis	¼ to 1 gr.
— Rhei	2 to 8 gr.
— Sarsæ Liquidum	2 to 4 fl. dr.

Name.	Dose.
Extractum Stramonii	$\frac{1}{4}$ to 1 gr.
— Strophanthi	$\frac{1}{4}$ to 1 gr.
— Taraxaci	5 to 15 gr.
— — Liquidum	$\frac{1}{2}$ to 2 fl. dr.
Fel Bovinum Purificatum	5 to 15 gr.
Ferri Arsenas	$\frac{1}{16}$ to $\frac{1}{4}$ gr.
— et Ammonii Citras...	5 to 10 gr.
— Carbonas Saccharatus	10 to 30 gr.
— Phosphas (71, 73)	5 to 10 gr.
— et Quininæ Citras	5 to 10 gr.
— Sulphas (58, 71, 73, 76)	1 to 5 gr.
— — Exsiccatus	$\frac{1}{2}$ to 3 gr.
Ferrum Redactum (73, 76)	1 to 5 gr.
— Tartaratum	5 to 10 gr.
Galbanum	5 to 15 gr.
Glusidum (60, 76)	$\frac{1}{2}$ to 2 gr.
Glycerinum (47, 54, 56, 57)	1 to 2 fl. dr.
— Pepsini	1 to 2 fl. dr.
Guaiaci Resina (34, 69)	5 to 15 gr.
Heroin Hydrochloridum (47) ...	$\frac{1}{12}$ to $\frac{1}{6}$ gr.
Hexamethylenetetramine (63) ...	5 to 15 gr.
Homatropinæ Hydrobromidum ...	$\frac{1}{80}$ to $\frac{1}{20}$ gr.
Hydrargyri Iodidum Rubrum... ...	$\frac{1}{32}$ to $\frac{1}{16}$ gr.
— Perchloridum	$\frac{1}{32}$ to $\frac{1}{16}$ gr.
Hydrargyrum cum Creta (72, 73, 74, 76)	1 to 5 gr.
Hyoscinæ Hydrobromidum	$\frac{1}{200}$ to $\frac{1}{100}$ gr.
Hyoscyaminæ Sulphas	$\frac{1}{200}$ to $\frac{1}{100}$ gr.
Infusum Aurantii (57)	$\frac{1}{2}$ to 1 fl. oz.
— — Compositum	$\frac{1}{2}$ to 1 fl. oz.
— Buchu (55)	1 to 2 fl. oz.
— Calumbæ (61)	$\frac{1}{2}$ to 1 fl. oz.

Name.	Dose.
Infusum Caryophylli	½ to 1 fl. oz.
— Cascarillæ	½ to 1 fl. oz.
— Chiratæ	½ to 1 fl. oz.
— Cinchonæ Acidum	½ to 1 fl. oz.
— Cuspariæ	1 to 2 fl. oz.
— Digitalis	2 to 4 fl. dr.
— Ergotæ	1 to 2 fl. oz.
— Gentianæ Compositum	½ to 1 fl. oz.
— Krameriæ	½ to 1 fl. oz.
— Lupuli	1 to 2 fl. oz.
— Quassiæ (57)	½ to 1 fl. oz.
— Rhei	½ to 1 fl. oz.
— Rosæ Acidum	½ to 1 fl. oz.
— Scoparii (63)	1 to 2 fl. oz.
— Senegæ (63)	½ to 1 fl. oz.
— Sennæ (43)	½ to 1 fl. oz.; as a draught, 2 fl. oz.
— Serpentariæ	½ to 1 fl. oz.
— Uvæ Ursi	½ to 1 fl. oz.
Injectio Apomorphinæ Hypodermica. ...	5 to 10 min. (By sub. injtn.)
— Cocainæ Hypodermica	2 to 5 min. (By sub. injtn.)
— Ergotæ Hypodermica	3 to 10 min. (By sub. injtn.)
— Morphinæ Hypodermica	2 to 5 min. (By sub. injtn.)
Iodoformum	½ to 3 gr.
Ipecacuanhæ Radix (73)	Expectorant, ¼ to 2 gr.; emetic, 15 to 30 gr.
Jalapa (72, 75)	5 to 20 gr.
Jalapæ Resina	2 to 5 gr.

Name.	Dose.
Kino	5 to 20 gr.
Liquor Ammonii Acetatis (57, 58, 62)	2 to 6 fl. dr.
— Citratis	2 to 6 fl. dr.
— Arsenicalis	2 to 8 min.
— Arsenici Hydrochloricus (58) ...	2 to 8 min.
— Arsenii et Hydrargyri Iodidi ...	5 to 20 min.
— Atropinæ Sulphatis	½ to 1 min.
— Bismuthi et Ammonii Citratis. ...	½ to 1 fl. dr.
— Calcis	1 to 4 fl. oz.
— — Saccharatus	20 to 60 min.
— Calumbæ Concentratus	½ to 1 fl. dr.
— Chiratæ Concentratus	½ to 1 fl. dr.
— Cuspariæ Concentratus	½ to 1 fl. dr.
— Ethyl Nitritis	20 to 60 min.
— Ferri Acetatis	5 to 15 min.
— — Perchloridi (57, 58)	5 to 15 min.
— — Pernitratis	5 to 15 min.
— Hydrargyri Perchloridi (59) ...	½ to 1 fl. dr.
— Hydrogenii Peroxidi	½ to 2 fl. dr.
— Krameriæ Concentratus	½ to 1 fl. dr.
— Magnesii Carbonatis (62)	1 to 2 fl. oz.
— Morphinæ Acetatis	10 to 60 min.
— — Hydrochloridi (47)	10 to 60 min.
— — Tartratis	10 to 60 min.
— Potassæ	10 to 30 min. freely diluted.
— Potassii Permanganatis	2 to 4 fl. dr.
— Quassiæ Concentratus	½ to 1 fl. dr.
— Rhei Concentratus	½ to 1 fl. dr.

Name.	Dose.
Liquor Sarsæ Compositus Concentratus. ...	2 to 8 fl. dr.
— Senegæ Concentratus	½ to 1 fl. dr.
— Sennæ Concentratus	½ to 1 fl. dr.
— Serpentariæ Concentratus... ...	½ to 2 fl. dr.
— Sodæ Chlorinatæ	10 to 20 min.
— Sodii Arsenatis	2 to 8 min.
— Strychninæ Hydrochloridi (58) ...	2 to 8 min.
— Thyroidei	5 to 15 min.
— Trinitrini	½ to 2 min.
Lithii Carbonas	2 to 5 gr.
— Citras	5 to 10 gr.
— — Effervescens	60 to 120 gr.
Lupulinum	2 to 5 gr.
Magnesia Levis	For repeated doses, 5 to 30 gr.; for a single dose, 30 to 60 gr.
— Ponderosa	
Magnesii Carbonas Levis (55, 60)	
— — Ponderosa (68, 69, 72) ...	
— Sulphas (43, 54, 56, 57, 59, 60, 62) ...	For repeated doses, 30 to 120 gr.; for a single dose, ¼ to ½ oz.
— — Effervescens	For repeated doses, 60 to 240 gr.; for a single dose, ½ to 1 oz.
Menthol (47)	½ to 2 gr.
Methyl Sulphonal (76)	10 to 30 gr.
Mistura Ammoniaci	½ to 1 fl. oz.
— Amygdalæ	½ to 1 fl. oz.
— Creosoti	½ to 1 fl. oz.

Name.	Dose.
Mistura Cretæ	½ to 1 fl. oz.
— Ferri Composita (57)	½ to 1 fl. oz.
— Guaiaci	½ to 1 fl. oz.
— Olei Ricini	1 to 2 fl. oz.
— Sennæ Composita	1 to 2 fl. oz. as a draught.
Morphinæ Acetas	⅛ to ½ gr.
— Hydrochloridum (76)	⅛ to ½ gr.
— Tartras	⅛ to ½ gr.
Naphthol	3 to 10 gr.
Nux Vomica	1 to 4 gr.
Oleum Anethi	½ to 3 min.
— Anisi (77)	½ to 3 min.
— Anthemidis	½ to 3 min.
— Cajuputi	½ to 3 min.
— Carui (62)	½ to 3 min.
— Caryophylli	½ to 3 min.
— Cinnamomi	½ to 3 min.
— Copaibæ	5 to 20 min.
— Coriandri	½ to 3 min.
— Crotonis	½ to 1 min.
— Cubebæ	5 to 20 min.
— Eucalypti	½ to 3 min.
— Juniperi	½ to 3 min.
— Lavandulæ	½ to 3 min.
— Limonis	½ to 3 min.
— Menthæ Piperitæ	½ to 3 min.
— — Viridis	½ to 3 min.
— Morrhuæ (37, 60)	1 to 4 fl. dr.
— Myristicæ	½ to 3 min.
— Phosphoratum	1 to 5 min.

Name.	Dose.
Oleum Pimentæ	$\frac{1}{2}$ to 3 min.
— Ricini (60)	1 to 8 fl. dr.
— Rosmarini	$\frac{1}{2}$ to 3 min.
— Santali (60)	5 to 30 min.
— Terebinthinæ	2 to 10 min.; as anthelmintic, 3 to 4 fl. dr.
Opium (72, 76)	$\frac{1}{2}$ to 2 gr.
Oxymel	1 to 2 fl. dr.
— Scillæ	$\frac{1}{2}$ to 1 fl. dr.
Paraldehydum	$\frac{1}{2}$ to 2 fl. dr.
Pepsinum (76)	5 to 10 gr.
Phenacetinum (74, 76)	5 to 10 gr.
Phenazonum (76)	5 to 20 gr.
Phenolphthalein (76)	$\frac{1}{2}$ to 4 gr.
Phosphorus	$\frac{1}{100}$ to $\frac{1}{20}$ gr.
Physostigminæ Sulphas	$\frac{1}{60}$ to $\frac{1}{20}$ gr.
Picrotoxinum	$\frac{1}{100}$ to $\frac{1}{25}$ gr.
Pilocarpinæ Nitras	$\frac{1}{20}$ to $\frac{1}{2}$ gr.
Pilula Aloes Barbadensis	4 to 8 gr.
— — et Asafetidæ	4 to 8 gr.
— — — Ferri	4 to 8 gr.
— — — Myrrhæ	4 to 8 gr.
— — Socotrinæ	4 to 8 gr.
— Cambogiæ Composita	4 to 8 gr.
— Colocynthidis Composita (72) ...	4 to 8 gr.
— — et Hyoscyami (73)	4 to 8 gr.
— Ferri (67)	5 to 15 gr.
— Galbani Composita	4 to 8 gr.
— Hydrargyri	4 to 8 gr.
— — Subchloridi Composita... ...	4 to 8 gr.

Name.	Dose.
Pilula Ipecacuanhæ cum Scilla ...	4 to 8 gr.
— Phosphori	1 to 2 gr.
— Plumbi cum Opio	2 to 4 gr.
— Quininæ Sulphatis	2 to 8 gr.
— Rhei Composita (74)	4 to 8 gr.
— Saponis Composita	2 to 4 gr.
— Scammonii Composita	4 to 8 gr.
— Scillæ Composita	4 to 8 gr.
Plumbi Acetas...	1 to 5 gr.
Podophylli Resina (76)	$\frac{1}{4}$ to 1 gr.
Potassii Acetas (63)	10 to 60 gr.
— Bicarbonas (60, 61)	5 to 30 gr.
— Bichromas	$\frac{1}{10}$ to $\frac{1}{5}$ gr.
— Bromidum (44, 56, 59, 61)	5 to 30 gr.
— Carbonas	5 to 20 gr.
— Chloras	5 to 15 gr.
— Citras (55, 57, 60, 63)	10 to 40 gr.
— Iodidum (59, 61)	5 to 20 gr.
— Nitras (34, 62)	5 to 20 gr.
— Permanganas	1 to 3 gr.
— Sulphas (70)	10 to 40 gr.
— Tartras	30 to 240 gr.
— — Acidus (68, 75)	20 to 60 gr.
Pulvis Antimonialis (75)	3 to 6 gr.
— Catechu Compositus	10 to 40 gr.
— Cinnamomi Compositus	10 to 40 gr.
— Cretæ Aromaticus (72)	10 to 60 gr.
— — — cum Opio (55)	10 to 40 gr.
— Elaterini Compositus	1 to 4 gr.
— Glycyrrhizæ Compositus	60 to 120 gr.
— Ipecacuanhæ Compositus (71, 74, 76)	5 to 15 gr.

Name.	Dose.
Pulvis Jalapæ Compositus	20 to 60 gr.
— Kino Compositus	5 to 20 gr.
— Opii Compositus	2 to 10 gr.
— Rhei Compositus	20 to 60 gr.
— Scammonii Compositus (75) ...	10 to 20 gr.
— Tragacanthæ Compositus	20 to 60 gr.
Quininæ Hydrochloridum	1 to 10 gr.
— — Acidum	1 to 10 gr.
— Sulphas (58, 61, 67, 72, 73, 76)	1 to 10 gr.
Rhei Radix (34, 59, 74)	3 to 10 gr. for repeated doses; for a single dose, 15 to 30 gr.
Salicinum (76)	5 to 20 gr.
Salol (76)	5 to 15 gr.
Santoninum (75)	2 to 5 gr.
Scammoniæ Resina (75)	3 to 8 gr.
Scammonium	5 to 10 gr.
Scilla (73)	1 to 3 gr.
Soda Tartarata...	120 to 240 gr.
Sodii Arsenas	$\frac{1}{40}$ to $\frac{1}{10}$ gr.
— Benzoas	5 to 30 gr.
— Bicarbonas (54, 56, 59, 68, 70, 74) ...	5 to 30 gr.
— Bromidum	5 to 30 gr.
— Carbonas	5 to 30 gr.
— — Exsiccatus	3 to 10 gr.
— Chloridum (70)	10 to 60 gr.; as an emetic, $\frac{1}{2}$ to 1 oz.
— Citro-Tartras Effervescens ...	60 to 120 gr.

Name.	Dose.
Sodii Hypophosphis	3 to 10 gr.
— Iodidum	5 to 20 gr.
— Nitris	1 to 2 gr.
— Phosphas (71)	30 to 120 gr. for repeated doses for a single dose $\frac{1}{4}$ to $\frac{1}{2}$ oz.
— — Acida (63)	30 to 60 gr.
— — Effervescens	60 to 120 gr. for repeated doses for a single dose $\frac{1}{4}$ to $\frac{1}{2}$ oz.
— Salicylas (63)	10 to 30 gr.
— Sulphas	30 to 120 gr. for repeated doses for a single dose $\frac{1}{4}$ to $\frac{1}{2}$ oz.
— — Effervescens	60 to 120 gr. for repeated doses for a single dose $\frac{1}{4}$ to $\frac{1}{2}$ oz.
— Sulphis	5 to 20 gr.
— Sulphocarbolas	5 to 15 gr.
Spiritus Ætheris (54, 55) ...	20 to 40 min. for repeated doses for a single dose 60 to 90 min.
— — Compositus	
— — Nitrosi (57, 62, 63) ...	
— Ammoniæ Aromaticus (54, 56, 61)	
— Ammoniæ Fetidus... ...	
— Anisi	5 to 20 min.
— Armoraciæ Compositus	1 to 2 fl. dr.
— Cajuputi	5 to 20 min.

Name.	Dose.
Spiritus Camphoræ (59)	5 to 20 min.
— Chloroformi (47)	5 to 20 min. for repeated doses; for a single dose, 30 to 40 min.
— Cinnamomi	5 to 20 min.
— Juniperi (63)	20 to 60 min.
— Lavandulæ	5 to 20 min.
— Menthæ Piperitæ	5 to 20 min.
— Myristicæ	5 to 20 min.
Strychnina (67, 73)	$\frac{1}{60}$ to $\frac{1}{15}$ gr.
Strychninæ Hydrochloridum	$\frac{1}{60}$ to $\frac{1}{15}$ gr.
Succus Belladonnæ	5 to 15 min.
— Conii	1 to 2 fl. dr.
— Hyoscyami (44)	$\frac{1}{2}$ to 1 fl. dr.
— Scoparii	1 to 2 fl. dr.
— Taraxaci	1 to 2 fl. dr.
Sulphonal (76)	10 to 30 gr.
Sulphur Precipitatum	20 to 60 gr.
— Sublimatum (34, 69)	20 to 60 gr.
Suprarenal Gland (76)	5 gr. & upwards.
Syrupus Aromaticus	$\frac{1}{2}$ to 1 fl. dr.
— Aurantii (60)	$\frac{1}{2}$ to 1 fl. dr.
— — Floris	$\frac{1}{2}$ to 1 fl. dr.
— Calcii Lactophosphatis	$\frac{1}{2}$ to 1 fl. dr.
— Cascaræ Aromaticus	$\frac{1}{2}$ to 2 fl. dr.
— Chloral	$\frac{1}{2}$ to 2 fl. dr.
— Codeinæ	$\frac{1}{2}$ to 2 fl. dr.
— Ferri Iodidi	$\frac{1}{2}$ to 1 fl. dr.
— — Phosphatis	$\frac{1}{2}$ to 1 fl. dr.
— — — cum Quinina et Strychnina ...	$\frac{1}{2}$ to 1 fl. dr.

Name.	Dose.
Syrupus Hemidesmi	½ to 1 fl. dr.
— Limonis (47)	½ to 1 fl. dr.
— Pruni Virginianæ	½ to 1 fl. dr.
— Rhei	½ to 2 fl. dr.
— Rhœados	½ to 1 fl. dr.
— Rosæ	½ to 1 fl. dr.
— Scillæ (47)	½ to 1 fl. dr.
— Sennæ (53)	½ to 2 fl. dr.
— Tolutanus (47)	½ to 1 fl. dr.
— Zingiberis (44, 57, 62)	½ to 1 fl. dr.
Tabellæ Trinitrini	1 to 2 tablets.
Terebenum	5 to 15 min.
Terpini Hydratum (47)	2 to 6 gr.
Theobrominum Sodio-Salicylicum (76)	5 to 15 gr.
Thymol	½ to 2 gr.
Thyroideum Siccum (76)	3 to 10 gr.
Tinctura Aconiti	5 to 15 min.; if very frequently repeated, 2 to 5 min.
— Aloes (53)	½ to 1 fl. dr. for repeated doses; for a single dose, 1½ to 2 fl. dr.
— Asafetidæ (54)	½ to 1 fl. dr.
— Aurantii (44)	½ to 1 fl. dr.
— Belladonnæ (59)	5 to 15 min.
— Benzoini Composita	½ to 1 fl. dr.
— Buchu	½ to 1 fl. dr.
— Calumbæ	½ to 1 fl. dr.
— Camphoræ Composita (47, 62) ...	½ to 1 fl. dr.

Name.	Dose.
Tinctura Cannabis Indicæ (44)	5 to 15 min.
— Cantharidis	5 to 15 min.; if frequently repeated, 2 to 5 min.
— Capsici (56)	5 to 15 min.
— Cardamomi Composita (53) ...	½ to 1 fl. dr.
— Cascarillæ	½ to 1 fl. dr.
— Catechu	½ to 1 fl. dr.
— Chiratæ	½ to 1 fl. dr.
— Chloroformi et Morphinæ Composita.	5 to 15 min.
— Cimicifugæ	½ to 1 fl. dr.
— Cinchonæ	½ to 1 fl. dr.
— — Composita	½ to 1 fl. dr.
— Cinnamomi	½ to 1 fl. dr.
— Cocci	5 to 15 min.
— Colchici Seminum	5 to 15 min.
— Conii	½ to 1 fl. dr.
— Croci	5 to 15 min.
— Cubebæ	½ to 1 fl. dr.
— Digitalis	5 to 15 min.
— Ergotæ Ammoniata	½ to 1 fl. dr.
— Ferri Perchloridi	5 to 15 min.
— Gelsemii	5 to 15 min.
— Gentianæ Composita	½ to 1 fl. dr.
— Guaiaci Ammoniata	½ to 1 fl. dr.
— Hamamelidis	½ to 1 fl. dr.
— Hydrastis	½ to 1 fl. dr.
— Hyoscyami (53, 55, 57, 59)	½ to 1 fl. dr.
— Iodi	2 to 5 min.

Name.	Dose.
Tinctura Jaborandi	½ to 1 fl. dr.
— Jalapæ	½ to 1 fl. dr.
— Kino	½ to 1 fl. dr.
— Krameriæ	½ to 1 fl. dr.
— Lavandulæ Composita	½ to 1 fl. dr.
— Limonis	½ to 1 fl. dr.
— Lobeliæ Ætherea (59)	5 to 15 min.
— Lupuli	½ to 1 fl. dr.
— Myrrhæ	½ to 1 fl. dr.
— Nucis Vomicæ (53, 57, 60, 61)	5 to 15 min.
— Opii (53, 55, 60)	5 to 15 min. for repeated doses; for a single dose, 20 to 30 min.
— — Ammoniata	½ to 1 fl. dr.
— Podophylli	5 to 15 min.
— Pruni Virginianæ	½ to 1 fl. dr.
— Quassiæ	½ to 1 fl. dr.
— Quillaiæ (35, 44)	½ to 1 fl. dr.
— Quininæ	½ to 1 fl. dr.
— — Ammoniata	½ to 1 fl. dr.
— Rhei Composita (54, 62)	½ to 1 fl. dr. for repeated doses; for a single dose, 2 to 4 fl. dr.
— Scillæ (63)	5 to 15 min.
— Senegæ	½ to 1 fl. dr.
— Sennæ Composita	½ to 1 fl. dr. for repeated doses; for a single dose, 2 to 4 fl. dr.

Name.	Dose.
Tinctura Serpentariæ	½ to 1 fl. dr.
-- Stramonii	5 to 15 min.
-- Strophanthi...	5 to 15 min.
— Sumbul	½ to 1 fl. dr.
— Tolutana	½ to 1 fl. dr.
— Valerianæ Ammoniata	½ to 1 fl. dr.
-- Zingiberis (53, 56, 60)	½ to 1 fl. dr.
Trional (76)	10 to 30 gr.
Veronal (75)	5 to 15 gr.
Vinum Antimoniale (62)	10 to 30 min.; as an emetic, 2 to 4 fl. dr.
— Colchici (57)	10 to 30 min.
— Ferri...	1 to 4 fl. dr.
— — Citratis	1 to 4 fl. dr.
— Ipecacuanhæ (54)...	10 to 30 min. as expectorant; as an emetic, 4 to 6 fl. dr.
-- Quininæ	½ to 1 fl. oz.
Zinci Acetas	1 to 2 gr.
-- Oxidum (75, 76)	3 to 10 gr.
-- Sulphas	1 to 3 gr. as a tonic; as an emetic, 10 to 30 gr.
— Valerianas (76)	1 to 3 gr.

NOTES.

NOTES.

NOTES.

NOTES.

NOTES.

NOTES.

NOTES.

NOTES.

NOTES.

℞ mist. Cyst. Co.

3 j. t. d. s. (3)

℞ mist. Magnes. Co.

3 j. t. d. s. (3)

℞ Pulv. Co. q. s.

Fiat mistura.

℞ Hyd. c. Creta pi.

Fiat pulvis.

℞. Bismuth. Carb. ℥x.
Sod. B. Carb. ℥xx
Tinct. Card. Co. 3 j.
Aq. m. Pip. ad 3 iv.
M. d. s. (4)

Ord. p. 2.

R. Pd. Bun. A x v.
Timothy Hayman & Son.
Agt. mark. Pipe 3
C. & S. (4).

---

on discharging.
Ry. Mist O. P. Dr
3 Jo.
2 5 - 6
(19)

Quantity of Hemp - 10. Days
24. 7.